Final Consolidated Atlantic Highly Migratory Species Fishery



Including:

A Final Environmental Impact Statement,
A Final Regulatory Impact Review,
A Final Regulatory Flexibility Analysis,
A Final Social Impact Assessment,
Framework Actions, and
the 2006 Stock Assessment and Fishery Evaluation Report



FINAL

CONSOLIDATED ATLANTIC HIGHLY MIGRATORY SPECIES FISHERY MANAGEMENT PLAN

Including:

A Final Environmental Impact Statement,
A Final Regulatory Impact Review,
A Final Regulatory Flexibility Analysis,
A Final Social Impact Analysis,
Framework Actions, and
the 2006 Stock Assessment and Fishery Evaluation Report

July 2006

Highly Migratory Species Management Division Office of Sustainable Fisheries National Marine Fisheries Service 1315 East-West Highway Silver Spring, Maryland 20910





Final Consolidated Atlantic Highly Migratory Species Fishery Management Plan

July 2006

Highly Migratory Species Management Division Office of Sustainable Fisheries National Marine Fisheries Service 1315 East-West Highway Silver Spring, Maryland 20910

The photos on the cover were contributed by the NOAA Photo Library And Mark Grace, SEFSC, NOAA Fisheries

Citation:

NMFS. 2006. Final Consolidated Atlantic Highly Migratory Species Fishery Management Plan. National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Office of Sustainable Fisheries, Highly Migratory Species Management Division, Silver Spring, MD. Public Document. pp. 1600.

The Final Consolidated Atlantic Highly Migratory Species Fishery Management Plan

Actions:

Consolidate the Fishery Management Plan for Atlantic Tunas, Swordfish, and Shark and the Atlantic Billfish Fishery Management Plan; establish workshops for fishermen and dealers; consider changes to time/area closures; address rebuilding and/or overfishing of northern albacore tuna, finetooth sharks, and Atlantic billfish; modify the management process of bluefin tuna; change the fishing year; modify the authorized gears; implement minor changes and clarifications to the regulations; and begin the process to update essential fish habitat

Type of Statement: Final Environmental Impact Statement; Final Regulatory Impact Review; Final Regulatory Flexibility Analysis; Final Social Impact Statement; and Final Framework Actions

Lead Agency:

National Marine Fisheries Service

For Further Information:

Karyl Brewster-Geisz

Highly Migratory Species Management Division F/SF1

1315 East-West Highway Silver Spring, MD 20910

(301) 713-2347; (301) 713-1917 (fax)

Abstract:

In 2003, the National Marine Fisheries Service (NMFS) began the process to amend the Fishery Management Plan for Atlantic Tunas, Swordfish, and Sharks and the Atlantic Billfish Fishery Management Plan. After considering comments on a scoping paper and on a predraft document, NMFS decided to consolidate these fishery management plans, modify the fishery management plan management measures as necessary, implement framework actions, and begin the process for updating essential fish habitat. The draft of this document was released on August 19, 2005. The comment period was open until March 1, 2006. During this time, 24 public hearings were held throughout the coastal states from Maine through Texas and the Caribbean. The final document describes a range of alternatives that could impact fishermen and dealers for all highly migratory species fisheries. The preferred alternatives include those to: establish mandatory workshops for fishermen and dealers; implement two small closures, consistent with regulations implemented by the Gulf of Mexico Fishery Management Council; establish criteria for modifying and/or establishing time/area closures; address rebuilding and/or overfishing of northern albacore tuna, finetooth sharks, and Atlantic billfish; modify the management process of bluefin tuna; change the fishing year for tunas, swordfish, and billfish back to a calendar year; authorize additional fishing gears; and clarify the regulations.

EXECUTIVE SUMMARY

Atlantic Highly Migratory Species (HMS) are managed under the dual authority of the Magnuson-Stevens Fishery Conservation and Management Act (Magnuson-Stevens Act) and the Atlantic Tunas Convention Act (ATCA). Under the Magnuson-Stevens Act, the National Marine Fisheries Service (NMFS) must manage fisheries to maintain optimum yield (OY) by rebuilding overfished fisheries and preventing overfishing. Under ATCA, NMFS is authorized to promulgate regulations, as may be necessary and appropriate, to implement the recommendations from the International Commission for the Conservation of Atlantic Tunas (ICCAT). Before this action, tunas, swordfish, and sharks were managed under the 1999 Fishery Management Plan (FMP) for Atlantic Tunas, Swordfish, and Sharks (and its 2003 amendment) and billfish were managed under the 1988 Atlantic Billfish FMP (and its 1999 amendment). This final HMS FMP combines the management of all Atlantic HMS into one FMP, and combines and simplifies the objectives of the previous FMPs.

NMFS announced its intent to prepare an Environmental Impact Statement to amend the two previous FMPs on July 9, 2003. In this notice, NMFS asked for comments on quota allocations of Atlantic bluefin tuna (BFT), swordfish, and sharks among and within domestic fishing categories; management alternatives to improve and streamline the current HMS limited access permit program; a review of HMS essential fish habitat (EFH) identifications; and exempted fishing and scientific research permitting issues. On April 30, 2004, NMFS announced the availability of an Issues and Options Paper and its intent to hold nine scoping meetings. This paper expanded the list of issues to include those issues listed above, additional issues for every species, HMS tournaments, bycatch reduction, recordkeeping and reporting, workshops, authorized fishing gears, and consolidation of the FMPs. NMFS presented the Issues and Options Paper to the New England, Mid-Atlantic, and Gulf of Mexico Fishery Management Councils and the Atlantic States Marine Fisheries Commission. A summary of the major comments received during scoping was released in December 2004 and is available on the HMS Management Division webpage at http://www.nmfs.noaa.gov/sfa/hms.

The Issues and Options paper included an exhaustive list of issues that NMFS could address regarding Atlantic HMS. During scoping, NMFS heard of more issues and options that merit additional consideration and examination. At the Predraft stage, in order to complete this action in a timely manner, NMFS decided to handle in this rulemaking only some of the issues identified in the Issues and Options paper and scoping process. NMFS prioritized the issues and chose to consider those that were required by law (*e.g.*, handling and release workshops are required under the 2004 Biological Opinion) and/or would improve the management of the fisheries (*e.g.*, amending the FMP for the BFT General Category should allow management to make changes in the fisheries on a more timely basis).

In February 2005, NMFS released the combined Predraft of the Consolidated HMS FMP and the 2005 Annual Stock Assessment and Fishery Evaluation (SAFE) Report. NMFS presented the Predraft document to all five Atlantic Fishery Management Councils, both the Gulf and Atlantic States Marine Fisheries Commissions, and to the HMS and Billfish Advisory Panels. Comments received on both the Issues and Options Paper and the Predraft were considered when drafting and analyzing the ecological, economic, and social impacts of the

alternatives in both the draft and final HMS FMP. A summary of the comments received on the Predraft was released in June 2005 and is available on the HMS Management Division webpage. While some of the options changed between the Predraft and Draft stages, the overall list of issues to be addressed did not change.

On August 19, 2005, the draft HMS FMP and proposed rule were released. Originally, the comment period was set to end 60 days after publication (October 18, 2005). However, due to hurricanes Katrina and Rita, NMFS extended the comment period to March 1, 2006 (for a total comment period of 194 days), in order to ensure that those fishermen directly affected by the hurricanes would have an adequate amount of time to review the document and provide comment. Several thousand written comments were received, 24 public hearings were held, and all five Atlantic Fishery Management Councils and the Gulf and Atlantic States Marine Fisheries Commissions were given briefings. A summary of the public comments received and NMFS' response to those comments is included in an appendix of this document and will also be in the final rule implementing the regulations. In addition to the public comments, NMFS also had three independent scientists (*i.e.*, scientists not involved in the drafting of the document) review three specific sections of the draft HMS FMP. The three sections were the time/area analyses, the standardized bycatch reporting methodology, and the review of EFH. The peer review comments are also included in an appendix of this document.

The preferred alternatives in this document considered all of the comments received from the general public at all stages of the rulemaking and the peer review by the independent scientists. Table 1 provides the list of the changes from the draft document and the expected implementation date of each alternative. A summary of the issues addressed and the other alternatives considered in this rulemaking can be found below. More detail can be found in Chapters 2 and 4 of this document. The final HMS FMP also consolidates the objectives for the FMP (listed in Chapter 1) and removes the exemption to the billfish no sale provision (allowed for, but not implemented, in the 1988 Billfish FMP). NMFS believes that the suite of preferred alternatives in this document should, consistent with the Magnuson-Stevens Act and other domestic laws, allow overfished Atlantic HMS to rebuild, address overfishing of Atlantic HMS, balance the needs of the fishermen and communities with the needs of the resource, and maximize OY for the fishery and the resource.

Table 1 The preferred alternatives at the draft and final stage of the Consolidated HMS FMP and the expected implementation date.

Preferred Alternative in Draft HMS FMP	Preferred Alternative in Final HMS FMP	Expected Implementation Date
Bycatch Reduction: Workshops		
A2. Mandatory workshops and certification for all HMS pelagic and bottom longline vessel owners	Same	January 1, 2007: must complete certification prior to renewing HMS permit in 2007
A3. Mandatory workshops and certification for vessel operators actively participating in HMS pelagic and bottom longline fisheries	Same	January 1, 2007: must complete certification prior to fishing on a vessel that has renewed its HMS permit in 2007
A5. Mandatory workshops and certification for shark gillnet vessel owners and operators	Same	January 1, 2007: must complete certification prior to renewing HMS permit in 2007
A6. Certification Renewal Timetable (Certification renewal every 3-years)	Same	30 days after final rule is published
A9. Mandatory HMS identification workshops for all shark dealers	Same	December 31, 2007
A16. Certification Renewal Timetable (Certification renewal every 3-years)	Same	30 days after final rule is published
Bycatch Reduction: Time/Area Closures		
B4. Implement complementary HMS management measures in Madison-Swanson and Steamboat Lumps Marine Reserves	Same	30 days after final rule is published
B5. Establish criteria to consider when implementing new time/area closures or making modifications to existing time/area closures	Same	30 days after final rule is published
Rebuilding and Preventing Overfishing: North	ern Albacore Tuna	
C3. Establish the foundation with ICCAT for developing an international rebuilding program	Same	30 days after final rule is published
Rebuilding and Preventing Overfishing: Fineto	ooth Sharks	
D4. Identify sources of finetooth shark fishing mortality to target appropriate management actions	Same	Ongoing
Rebuilding and Preventing Overfishing: Atlant	ic Billfish	
E3. Effective January 1, 2007, limit all Atlantic billfish tournament participants to using only non-offset circle hooks when using natural baits or natural bait/artificial lure combinations	E3. Effective January 1, 2007, limit all HMS permitted vessels participating in Atlantic billfish tournaments to deploying only non-offset circle hooks when using natural baits or natural bait/artificial lure combinations	January 1, 2007

Preferred Alternative in Draft HMS FMP	Preferred Alternative in Final HMS FMP	Expected Implementation Date
E6. Effective January 1, 2007, implement ICCAT Recommendations on Recreational Marlin Landings Limits	Same	January 1, 2007
E7. Effective January 1, 2007 - December 31, 2011, allow only catch and release fishing for Atlantic white marlin	No longer preferred	NA
Management Program Structure: Bluefin Tuna	Quota Management	
F3. Amend the management procedures regarding General category time-periods, subquota, as well as geographic set-asides to allow for future adjustments to take place via a regulatory framework action	Same	30 days after final rule is published
F3(c). Revise General category time-periods and subquotas to allow for a formalized winter fishery (June-Aug, 50%; Sept, 26.5%; Oct-Nov, 13%; Dec, 5.2% and Jan, 5.3%)	Same	30 days after final rule is published
F4. Clarify the procedures for calculating the Angling category school size-class BFT subquota allocation and remove the Angling category north/south dividing line	F4. Clarify the procedures for calculating the Angling category school size-class BFT subquota allocation and maintain the Angling category north/south dividing line	30 days after final rule is published
F6. Revise the annual BFT specification process to refer back to the supporting analytical documents of the Consolidated HMS FMP and include seasonal management measures in annual framework actions	Same	30 days after final rule is published
F8. Establish an individual quota category carry-over limit of 100 percent of the baseline allocation (<i>i.e.</i> , no more than the annual baseline allocation may be carried forward), except for the Reserve category, and authorize the transfer of quota exceeding the 100 percent limit to the Reserve or another domestic quota category, while maintaining status quo overharvest provisions	Same	30 days after final rule is published
F10. Revise and consolidate criteria considered prior to performing inseason and some annual BFT management actions	Same	30 days after final rule is published
Management Program Structure: Timeframe fo	or Annual Management of HMS Fi	sheries
G2. Shift the fishing year to January 1 – December 31 for all HMS	Same	January 1, 2008
Management Program Structure: Authorized F	ishing Gears	
H2. Authorize speargun fishing gear as a permissible gear type in the recreational Atlantic tuna fishery	H2. Authorize speargun fishing gear as a permissible gear type in the recreational Atlantic BAYS tuna fishery	30 days after final rule is published

Preferred Alternative in Draft HMS FMP	Preferred Alternative in Final HMS FMP	Expected Implementation Date
H4. Authorize green-stick for the commercial harvest of Atlantic BAYS tunas	No longer preferred	NA
H5. Authorize buoy gear in the commercial swordfish handgear fishery, and limit vessels employing buoy gear to possessing and deploying no more than 35 individual buoys, with each having no more than two hooks or gangions attached	H5. Authorize buoy gear as a permissible gear type in the commercial swordfish handgear fishery; limit vessels employing buoy gear to possessing and deploying no more than 35 floatation devices, with each individual gear having no more than two hooks or gangions attached	30 days after final rule is published
H7. Clarify the allowance of hand-held cockpit gears used at boat side for subduing HMS captured on authorized gears	Same	30 days after final rule is published
Management Program Structure: Regulatory H	lousekeeping	
I1(b). Establish additional restrictions on longline gear in HMS time/area closures by specifying a maximum and minimum allowable number of commercial fishing floats to qualify as a BLL and PLL vessel, respectively	No longer preferred	NA
I1(c). Differentiate between PLL and BLL gear based upon the species composition of the catch onboard or landed	Same	30 days after final rule is published
I2(b). Require that the 2 nd dorsal fin and the anal fin remain on all sharks through landing	Same	30 days after final rule is published
I3(b). Add new prohibition at § 635.71(a)(48) making it illegal for any person to, "Purchase any HMS that was offloaded from an individual vessel in excess of the retention limits specified in §§ 635.23 and 635.24"	Same	30 days after final rule is published
I3(c). Add new prohibition at § 635.71(a)(49) making it illegal for any person to, "Sell any HMS that was offloaded from an individual vessel in excess of the retention limits specified in §§ 635.23 and 635.24"	Same	30 days after final rule is published
I4(b). Amend the second coordinate of the East Florida Coast closed area so that it corresponds with the EEZ	Same	30 days after final rule is published
I5(b). Amend the definition of "handline" at § 635.2 by requiring that they be attached to, or in contact with, all vessels	Same	30 days after final rule is published
I6(b). Prohibit vessels issued commercial permits and operating outside of a tournament from possessing, retaining, or taking Atlantic billfish from the management unit	Same	30 days after final rule is published

Preferred Alternative in Draft HMS FMP	Preferred Alternative in Final HMS FMP	Expected Implementation Date
I7(b). Amend the HMS regulations to provide an option for Atlantic tunas dealers to submit required BFT reports using the Internet	Same	30 days after final rule is published
I8(b). Require submission of "No Fishing" reporting forms for selected vessels if no fishing trips occurred during the preceding month, postmarked no later than seven days after the end of the month	Same	30 days after final rule is published
I8(c). Require submission of the trip "Cost- Earnings" reporting form for selected vessels 30 days after a trip and the annual "Cost- Earning" report form by January 31 of each year	I8(c). Require submission of the trip "cost-earnings" reporting form for selected vessels 30 days after a trip, and the "annual expenditures" report form by the date specified on the form	30 days after final rule is published
I9(b). Require vessel owners to report non- tournament recreational landings of North Atlantic swordfish and Atlantic billfish	I9(b). Require vessel owners (or their designees) to report non-tournament recreational landings of North Atlantic swordfish and Atlantic billfish	30 days after final rule is published
I10(b). Modify the HMS regulations to state that "In addition, each year, 25 mt (ww) will be allocated for incidental catch by pelagic longlines" in the NED	I10(c). Conduct additional discussions at ICCAT regarding quota rollovers and adjust quotas allocated to account for bycatch related to pelagic longline fisheries in the vicinity of the management area boundary accordingly	30 days after final rule is published
I11(b). Require recreational vessels with a Federal permit to abide by Federal regulations, regardless of where they are fishing, unless a state has more restrictive regulations	Same	30 days after final rule is published

Bycatch Reduction: Workshops

The June 2004 Biological Opinion (BiOp) for the Atlantic HMS pelagic longline fishery requires NMFS to conduct training workshops regarding the safe release and disentanglement of sea turtles from pelagic longline gear and to certify that fishermen have attended these workshops. The October 2003 BiOp on the Atlantic shark fishery requires a series of workshops that provide gear handling techniques and protocols that deal with entanglements and protected species, in general, and include information on smalltooth sawfish and HMS requirements. Additionally, in Amendment 1 to the Atlantic Tunas, Swordfish, and Shark FMP, NMFS stated that if shark fishermen can show that they can fish for specific species (*e.g.*, target sandbar sharks) and correctly identify the shark species caught on their gear, then the Agency might consider using species-specific shark quotas in the future. Thus, NMFS felt it was important to

consider workshops, particularly workshops for handling and release of protected species and workshops for identification of Atlantic HMS, in this rulemaking.

The workshops for the safe release, disentanglement, and identification of protected resources are designed to reduce the post-hooking mortality of sea turtles and other protected resources by educating fishermen on how to apply the appropriate safe handling and release protocols, improve compliance with regulations, and enhance the utility of vessel logbook data. The preferred alternatives for the protected species workshops would require all longline and gillnet permit holders and operators to attend and be certified in handling and release techniques and gear. Mandatory workshops for vessel owners would be linked to the vessels' permit, ensuring well attended workshops. Including operators would guarantee at least one person on board the vessel during fishing activities is adept at the safe handling and release protocols. NMFS also considered a range of alternatives for the protected species workshops including voluntary workshops (no action) and mandatory workshops for the owners, operators, and the crew of all HMS longline and gillnet vessels.

The preferred alternative for the identification workshops calls for all Federally permitted shark dealers, or a designated proxy, to attend one-day workshops on species-specific identification of offloaded shark carcasses. NMFS believes that identifying shark carcasses is more difficult and uncertain than identifying other HMS carcasses as evidenced by the large proportion of "unclassified" sharks listed on shark dealer logbooks. This uncertainty compromises quota monitoring and stock assessment efforts. Dealers are a focal point for gathering shark landings information as sharks from numerous vessels are offloaded at each individual dealer. Positive identification is often less difficult for fishermen than dealers as they know exactly where (depth, type of habitat, etc) a shark has been caught and often see the sharks alive and intact. NMFS considered a range of alternatives for these identification workshops including voluntary HMS identification workshops for dealers, recreational fishermen, and all commercial vessel owners and operators (no action) and mandatory identification workshops for all HMS dealers and/or HMS permit holders.

Under the preferred alternatives, longline and gillnet permit holders and vessel operators and shark dealers would be required to be recertified every three years. NMFS also considered recertification time periods of two and five years. Requiring recertification every three years would balance the ecological benefits of maintaining familiarity with the protocols and the economic impacts of workshop attendance due to travel costs and lost fishing opportunities.

None of the preferred alternatives changed significantly between the draft and final stages of this HMS FMP, although NMFS did adjust the effective dates as a result of public comment and the lengthening of the comment period. These one-day workshops are not expected to result in excessive economic impacts as they would be scheduled at numerous locales along the Atlantic coast, minimizing travel and lost fishing time.

Bycatch Reduction: Time/Area Closures

Since the 1999 FMP for Atlantic Tunas, Swordfish, and Sharks, NMFS has implemented a number of time/area closures in order to reduce bycatch, to the extent practicable, consistent

with the National Standards. While the results of preliminary analyses examining the efficacy of these closures have been included in annual SAFE Reports, a comprehensive analysis of the impact of the closures on bycatch rates, the fishermen, and the communities is contained in this document. In this document, NMFS examines the current time/area closures to determine if these closures are accomplishing the original goals of the closures and whether changes are needed to accomplish other objectives. The results of that examination indicate that both bycatch and overall effort in the fleet has been reduced (see discussions of alternative B1 in Chapter 4).

In this HMS FMP, NMFS is preferring two alternatives in regard to time/area closures. The first preferred alternative would establish HMS regulations in the Madison-Swanson and Steamboat Lumps Marine Reserves that complement the Gulf of Mexico Fishery Management Council's regulations. These closures are expected to have minimal ecological, economic, or social impacts on HMS fishermen. The second preferred alternative would establish criteria that would guide future decision-making regarding implementation or modification of time/area closures. This would provide enhanced transparency, predictability, and understanding of HMS management decisions, allow for more adaptive management, and should result in minimal social and economic impacts. Any impacts for specific closures would be analyzed when those closures are considered.

As described in Chapter 4 and Appendix A, NMFS used POP and HMS logbook data to identify new areas for time/area closures and selected alternatives based on these data to further analyze 10 different closures or modifications for this rulemaking. NMFS evaluated the reduction in discards of white marlin, blue marlin, sailfish, spearfish, leatherback sea turtles, loggerhead sea turtles, other sea turtles, and BFT without redistribution of effort based on POP data and the HMS logbook data for the various time/area closure alternatives (see Chapter 4). Using HMS logbook data (see Chapter 4 and Appendix A), NMFS evaluated different scenarios of a redistribution of fishing effort model, where each scenario had different assumptions regarding how fishing effort would be redistributed into open areas. The model used in this time/area analysis was consistent with the methods used in past rulemakings (for more information on redistribution of effort model selection, please see page 4-6). Additional redistribution of effort scenarios were considered based on comments received on the Draft Consolidated HMS FMP and the OMB reviews. As described in Chapter 4, each scenario of the models had different assumptions regarding how fishermen would react to the closures (e.g., will fishermen move out of the closed area but continue fishing in surrounding open areas, move their business, or sell their permits to someone near an open area). Because of the difficulty in predicting fishermen's behavior, NMFS analyzed the range of what would happen fleet-wide while recognizing that individuals within the fleet may act differently, and large closures may result in more movement in order for fishermen to find open areas to fish and stay in business.

NMFS examined a wide range of alternatives including closing additional closures or combining these additional closures for pelagic longline gear in the Gulf of Mexico and the Atlantic Ocean, modifying existing closures for pelagic longline gear, establishing a closure for bottom longline gear to protect smalltooth sawfish, and closing all areas to pelagic longline gear. These alternatives were not preferred for a variety of reasons. The ecological benefits of some of the additional closure alternatives considered were predicted to be variable with redistribution of

effort, with potential negative ecological impacts to several species. For example, alternative B2(a) (May - Nov), intended primarily to reduce leatherback sea turtle interactions, and white marlin and BFT discards, could result in a 7.9 percent increase in loggerhead sea turtle interactions and a 10.3 percent increase in BFT discards (see Table 4.2). As described in Appendix A, even the modified redistribution of effort model for alternative B2(a) predicted increases in sailfish discards (4.7 percent), LCS discards (4.4 percent), BFT discards (1.6 percent), and BAYS discards (0.7 percent). When closure areas were combined, the redistribution of effort model predicted similar results with an increase in discards of several species.

Alternatives B3(a) and B3(b) were considered to refine existing closures and to provide additional opportunity to harvest legal-sized swordfish while not increasing bycatch. NMFS, however, is not preferring any modifications to the current closures. None of the modifications considered would have resulted in a large enough increase in retained catch to alleviate concerns over uncaught portions of the swordfish and BFT quotas. For instance, B3(a) was predicted to increase retained swordfish catch by only 30.72 mt, and B3(a) was predicted to increase the retained swordfish catch by 0.07 mt. However, as of April 30, 2006, 4,905.9 mt and 294.7 mt of directed and incidental quota, respectively, were still available for the 2005 fishing year. In addition, modifications to existing closures could result in increased bycatch of blue and white marlin, which is a concern given the stock status of blue and white marlin and the scheduled white marlin ESA review. Increased interactions with sea turtles and marine mammals (*e.g.*, pilot whales and Risso's dolphins) are an additional concern.

Finally, all of these analyses (those analyzing the impacts of new closures and those analyzing the impacts of modifications to existing closures) were conducted using J-hook data. New circle hook management measures were put into place in 2004, and NMFS is still assessing the effects of circle hooks on bycatch rates for HMS. Based on the Northeast Distant experiment, circle hooks likely have a significantly different catch rate than J-hooks. Therefore, NMFS needs to conduct further investigations to determine the potential impact of any new time/area closures or modifications to existing closures. NMFS anticipates that 2005 HMS logbook final data will become available in the summer of 2006. In addition, NMFS is awaiting additional information regarding the status of the pelagic longline fleet after the devastating hurricanes in the Gulf of Mexico during the fall of 2005. A majority of the pelagic longline fleet was thought to be severely damaged or destroyed during the 2005 hurricane season. The amount of pelagic longline fishing effort, especially within the Gulf of Mexico, will likely be assessed in the summer of 2006 when 2005 HMS logbook final data becomes available. Until NMFS can better estimate the current fishing effort and potential recovery of the pelagic longline fleet, it may be premature to implement any new time/area closures, particularly in the Gulf of Mexico. Furthermore, a number of stock assessments will be conducted during 2006 (blue marlin, white marlin, north and south swordfish, eastern and western BFT, and large coastal sharks). NMFS is waiting on the results of these stock assessments to help determine domestic measures with regard to management of these species.

For the bottom longline closure alternative (B6), NMFS is waiting for the Smalltooth Sawfish Recovery Team to designate critical habitat in order to compare possible closure areas with the critical habitat. Closing all areas to pelagic longline gear (alternative B7) would have

severe economic and social impacts in the short term and possible negative ecological impacts in the long term if U.S. quotas are transferred to countries without the same conservation ethic.

While NMFS did not change the preferred alternatives between the draft and final stages, NMFS did conduct additional analyses as a result of public comment. These analyses include examining the redistribution of effort model and its applicability, the mobility of the fleet, and the concept of a decision matrix. NMFS also began looking at the 2004 circle hook data for the pelagic longline fishery. In the future, NMFS intends, among other things, to investigate the choices fishermen have made regarding previous closures and to pursue alternatives to reduce bycatch in the Gulf of Mexico, especially for BFT. For BFT, NMFS is currently trying to assess how protecting one age class at the potential detriment of other age classes will affect the fish stock as a whole, and is also considering developing incentives that would dissuade fishermen from keeping incidentally caught BFT, particularly spawning BFT, in the Gulf of Mexico. This may involve research on how changes in fishing practices may help reduce bycatch of non-target species as well as the tracking of discards (dead and alive) by all gear types. More information on these additional analyses, their results, and potential future actions are contained in Chapter 4 and Appendix A.

Rebuilding and Preventing Overfishing: Northern Albacore Tuna

Since the 1999 FMP, NMFS has determined that northern albacore tuna are overfished. While NMFS published a final rule that stated NMFS would work with ICCAT to rebuild northern albacore, a rebuilding plan was not previously incorporated in the FMP. The preferred alternative would establish a foundation with ICCAT for developing an international rebuilding plan. Under this alternative, NMFS will continue to work with ICCAT member nations to develop and adopt an appropriate international rebuilding plan for northern albacore tuna with a specified recovery period, biomass targets, fishing mortality rate limits, and explicit interim milestones. The U.S. harvest of the North Atlantic stock is proportionally so low that the socioeconomic impacts to the United States would likely be minimal but would depend upon the specifics of the rebuilding plan adopted by ICCAT. The other alternatives of no action or unilateral action would not be consistent with the Magnuson-Stevens Act or ATCA, and would be unlikely to rebuild northern albacore.

Rebuilding and Preventing Overfishing: Finetooth Sharks

In 2002, NMFS determined that overfishing is occurring on finetooth sharks. In the 2003 Amendment to the 1999 FMP, because most finetooth landings appear to come from fishermen in non-HMS fisheries, NMFS stated that it would take action to identify sources of fishing mortality on finetooth sharks, increase outreach, improve enforcement of the recreational limits, and work with the Regional Fishery Management Councils to identify fisheries that catch finetooth sharks.

In this HMS FMP, NMFS prefers an alternative that would establish a plan to prevent overfishing. This preferred alternative would identify the sources of fishing mortality for finetooth sharks. The analyses in the HMS FMP found that the majority of finetooth sharks are landed in the South Atlantic region (primarily Florida) by vessels deploying gillnet gear and in possession of both a Spanish mackerel permit and a commercial shark permit. NMFS also found

that an unmanaged fishery, the southern kingfish fishery, also catches finetooth sharks. Thus, any management measures that are solely directed at fishermen using gillnet gear and in possession of a commercial shark permit could easily be circumvented by fishermen using gillnets for Spanish mackerel or kingfish. In addition to conducting analyses, NMFS has also contacted the states and Regional Fishery Management Councils, sent a letter to the South Atlantic Fishery Management Council requesting collaboration in management between gillnet fisheries, and requested that finetooth sharks be added to observer programs such as the Gulf of Mexico shrimp trawl fishery. These actions should provide additional options to address this issue.

NMFS considered other alternatives including no action, management measures targeting commercial shark permit holders, and management measures targeting recreational HMS permit holders. Targeting commercial shark permit holders is confounded by the fact that finetooth sharks are within the SCS complex, which is not currently overfished or experiencing overfishing, and commercial fishermen have only caught, on average, 28.5 percent of the SCS quota between 1999-2003. Measures aimed at the recreational fishery would only affect a small portion of the overall finetooth shark landings. Furthermore, a conservative bag limit of one shark (including finetooth shark) and a minimum size above the age at first maturity for males and females are already in place. NMFS intends to conduct a new small coastal shark stock assessment following the Southeast Assessment, Data, and Review process starting in 2007. As more research and data become available, NMFS may reconsider these other alternatives.

NMFS did not change the preferred alternative between the draft and final stages. NMFS believes that the preferred approach constitutes a plan to prevent overfishing and is a prudent means of establishing regulations that might affect a type of gear (gillnet), rather than an individual permit. Applying the regulations to the gear is critical as regulations implemented only on shark permit holders would only affect a sub-set of the individuals responsible for finetooth shark fishing mortality, could be easily circumvented, and would likely result in additional dead discards of finetooth sharks.

Rebuilding and Preventing Overfishing: Atlantic Billfish

Despite the implementation of domestic and international management measures, the status of Atlantic blue and white marlin has continued to decline. Currently, the status of sailfish and spearfish is uncertain. Atlantic white marlin has been identified as one of the most severely overfished species of any stock under ICCAT's purview for the past four years, but nevertheless continues to be subjected to unsustainable levels of fishing mortality throughout the Atlantic. In 2002, the United States undertook a status review of white marlin pursuant to the Endangered Species Act (ESA). While the status review team determined that white marlin stock status did not warrant a listing at that time, it concluded that "unless fishing mortality is reduced significantly and relatively quickly, the stock could decline to a level that would warrant ESA protection" (White Marlin Status Review Team 2002). NMFS will conduct another ESA listing review in 2007. As such, in this document, NMFS reviewed the current data and examined methods of reducing billfish mortality in both the commercial (*e.g.*, time/area closures) and recreational fisheries (*e.g.*, circle hook requirements).

NMFS is preferring two alternatives to reduce the post-release mortality of billfish associated with the directed billfish fishery. The first preferred alternative would require the use of non-offset circle hooks by HMS permitted vessels in billfish tournaments when using natural baits or natural bait/artificial lure combinations. The second preferred alternative would codify the ICCAT landings recommendations for billfish. The current landings recommendation would limit the United States to landing no more than 250 blue or white marlin per year. These alternatives strike a balance between conserving living marine resources and maintaining robust recreational fisheries while achieving the objectives of the HMS FMP. The preferred alternatives are anticipated to substantially reduce the post-release mortality of Atlantic white marlin, provide positive ecological benefits for other species such as blue marlin, sailfish, and tunas, and maintain consistency with United States' international obligations. NMFS is delaying the effective date for the circle hook requirement to mitigate, to the extent practicable, adverse economic impacts and losses in angler consumer surplus by allowing: tournament operators adequate time to adjust advertising, rules, business practices, and tournament formats; existing stockpiles of J-hooks to be used; and, anglers time to become comfortable and proficient with newly required gear.

As a result of public comment, NMFS is no longer preferring the alternative that would prohibit the landing of white marlin. Additionally, NMFS clarified the intent of the first preferred alternative to ensure that only HMS permit holders, not all tournament participants, would be affected by the circle hook requirement.

Management Program Structure: Bluefin Tuna Quota Management

Western Atlantic BFT are overfished, and one of the main objectives of the Consolidated HMS FMP is to end overfishing and rebuild overfished stocks, while providing reasonable fishing opportunities to harvest the limited quota that is available under the BFT rebuilding plan. Since the 1999 FMP, BFT management has become increasingly complicated and difficult for the public to understand and may no longer accurately reflect the needs of the fishery and goals of the 1999 FMP. These issues are evident on a daily basis from the number of constituent inquiries addressed by NMFS and the number of inseason management actions necessary throughout the season. In addition, NMFS has received a petition from the State of North Carolina Department of Marine Fisheries (NMDMF) for rulemaking to adjust the quota allocations to provide for a General category fishery off North Carolina in the winter. NMFS considers these requests and considers ways of clarifying BFT management.

Two of the preferred alternatives would amend the time period and sub quotas for the General category and clarify the procedures for calculating the Angling category school-size fish. These alternatives are expected to enhance NMFS' flexibility to address inherent variability in the BFT fishery while still allowing for business planning. They also respond in part to the NCDMF's Petition for Rulemaking and would allow for a formal General category winter BFT fishery while still recognizing the historical BFT catch rates in the New England area fishery. These preferred alternatives would also clarify the procedures NMFS used to implement the ICCAT recommendation regarding the eight percent tolerance limit of school BFT as well as maintain the recreational North/South dividing line as a management tool.

Two other preferred alternatives would provide participants in the BFT fishery a timely and stable baseline quota allocation from one year to the next, the ability to address under/overharvest from the previous year, the ability to establish the General category effort controls as well as recreational and commercial handgear daily retention limits for the upcoming season, and streamline the annual rulemaking process. Additionally, providing NMFS the authority to implement a cap on the amount of quota that may be carried forward from one fishing year to the next would allow NMFS to manage to harvest of BFT with more finite precision and minimize the occurrence of 'stockpiling' in any one quota category.

Another preferred alternative would consolidate and refine the criteria that NMFS must consider prior to conducting any inseason, and some annual, actions. This preferred alternative would assist in meeting the Consolidated HMS FMP's objectives in a consistent manner, providing reasonable fishing opportunities, increasing the transparency in the decision making process, and balancing the resource's needs with users' needs.

Management Program Structure: Timeframe for Annual Management of HMS Fisheries

In the 1999 FMP and 1999 Billfish Amendment, NMFS established a fishing year management cycle for tunas, billfish, and swordfish that began on June 1 and went through the following May 31. This fishing year was established to allow NMFS time to implement recommendations from ICCAT before the fishing year began. The change to the fishing year, however, has been problematic given that many of the data infrastructure and reporting requirements both within NMFS and ICCAT are based on a calendar year rather than a fishing year. NMFS prefers the alternative that would establish a fishing year management cycle for all HMS of January 1 through December 31. This preferred alternative is expected to simplify the regulatory process for constituents in the long term by managing all HMS fisheries on a calendar year and improve the United States' basis for negotiation at international forums.

Management Program Structure: Authorized Fishing Gears

In 1999, NMFS published a list of authorized gears for all fisheries across the nation. Occasionally, NMFS receives requests to modify the list of authorized gears. Sometimes, these requests include gear that fishermen use in other oceans or elsewhere in the Atlantic to catch the same species; other times, the requests are due to additional groups requesting to use a gear that is approved for one permit, but not another. NMFS considers some of these requests (*e.g.*, green-stick gear and speargun fishing gear) pertaining to HMS in this rulemaking.

NMFS prefers several alternatives that would add authorized gear types in HMS fisheries. The first preferred alternative would allow spearfishermen to participate in the Atlantic bigeye, albacore, yellowfin, and skipjack (BAYS) tunas fishery. This alternative is responsive to specific public comment and requests from constituents. This preferred alternative is anticipated to result in minimal negative ecological impacts and positive social and economic benefits. This preferred alternative is modified slightly from what was proposed in that, due to concerns related to the status of BFT, only BAYS tunas could be taken by spearfishermen, not BFT.

The second preferred alternative would allow the commercial swordfish handgear fishery to continue to utilize individual unattached buoyed gears (a.k.a. buoy gear), and would limit the maximum number of gears deployed by a vessel. Before this FMP, both recreational and commercial swordfish handgear fishermen could use this gear, previously called handline, and were not limited in the number of gears that could be deployed. This alternative may provide some positive ecological benefits by limiting future expansion of this gear sector and possibly by reducing the amount of lost fishing gear. This alternative could result in positive social benefits and would maintain current economic benefits to this sector. The last preferred alternative would, in response to requests from fishery participants, clarify the allowable use of secondary cockpit gears. This alternative should not result in an increase in bycatch mortality, over current levels, as secondary gears are currently utilized in HMS fisheries.

Although NMFS originally preferred an alternative that would allow for the use of greenstick in the commercial BAYS fishery in the Draft HMS FMP, it is not preferred in the Final HMS FMP. During the comment period, NMFS realized that many fishermen, both commercial and recreational, did not understand which gear configurations were currently allowed and which configurations the Agency was proposing to allow. Thus, NMFS will clarify the existing regulatory regime and the allowable configurations of green-stick gear in an effort to reduce confusion regarding the authorized use of green-stick gear.

Management Program Structure: Regulatory Housekeeping

This rulemaking also considers a number of corrections and additions to the Atlantic HMS regulations at 50 CFR part 635 and other relevant sections in the CFR (*e.g.*, 50 CFR part 300 contains information regarding international trade) in order to clarify their intent, remove incorrect cross-references, remove dated regulations, as appropriate, and aid enforcement. Besides the more than 40 minor corrections to the regulatory text, NMFS also considered a few changes that required alternatives. In all, NMFS is preferring 13 alternatives in this section across a wide range of eleven different issues.

The first issue in this section pertains to the definitions of bottom and pelagic longline gear. These gears catch different species and are currently differentiated by the number of weights and/or floats each gear uses. This raises enforcement concerns particularly in closed areas. As such, NMFS is preferring an alternative that would differentiate between gears based upon the species composition of the catch onboard or offloaded. This alternative is expected to accommodate the majority of commercial fishing operations, yet still provide a quantifiable method to differentiate between vessels using one gear or the other. Vessels that fish mixed trips (*i.e.*, trips that use both gear types) could still transit the closed areas provided the signals from their vessel monitoring system unit indicate the vessel is transiting and not fishing. This alternative is not expected to create significant adverse economic and social impacts and is expected to improve the monitoring of, and compliance with, HMS closed area regulations. NMFS originally preferred both the current preferred alternative and an alternative that would limit the number of floats on bottom longline vessels. NMFS is no longer preferring that alternative based upon public comment regarding impacts to vessel's operational flexibility, difficulties with terminology, and impracticalities in enforcing the alternative. Other alternatives

considered, besides the no action, included requiring time and depth recorders and closing all areas to "longline" rather than trying to define the gears.

The second issue pertains to shark identification. Currently, shark fishermen may remove all fins from the shark, consistent with the five-percent shark fin ratio. NMFS prefers an alternative that would require the second dorsal and anal fins to remain on all sharks through the first port of landing. While this alternative could have some minor economic and social impacts, this alterative is expected to generate ecological benefits by enhancing and improving species identification and data collection, thereby leading to improved management and increased shark populations. NMFS also considered alternatives that would allow fishermen to remove the second dorsal and anal fins from some species (*e.g.*, lemon sharks) or require all fins to remain on the shark.

In a third issue regarding sales of illegal landings, NMFS is preferring two alternatives that would add clear prohibitions to the regulations regarding the sale and purchase of landings in excess of the commercial retention limits. These alternatives may act as an additional deterrent to discourage this illegal practice. NMFS believes that the social benefits of preventing this practice should outweigh any short-term economic benefit gained as a result of illegally selling catches in excess of the commercial retention limits.

In a fourth issue regarding the definition of the closed areas, NMFS is preferring an alternative that would amend the area of the East Florida Coast closed area by extending one of its coordinates 1.02 km (0.55 nmi) seaward so that it corresponds with the outer boundary of the exclusive economic zone (EEZ). This alternative is not expected to create significant adverse economic and social impacts. Any fishing effort that would have occurred in this area would likely relocate to nearby open areas with similar catch rates. Because the East Florida Coast closed area would be enlarged under this alternative, it could reduce the bycatch of undersized swordfish, sailfish, and other HMS as compared with the no action alternative, but this reduction is expected to be minimal.

The fifth issue pertains to the definition of handline. In the authorized fishing gear section of the HMS FMP, NMFS is preferring an alternative that would define unattached handlines as buoy gear and restrict their use to commercial swordfish fishermen. In this section, NMFS is preferring an alternative that would require that handlines remain attached to all vessels. This alternative would primarily affect recreational fishery participants and commercial permittees that do not possess a commercial swordfish handgear permit. This alternative is not expected to have significant adverse social or economic impacts on fishery participants.

The sixth issue described in this section pertains to the retention of billfish by commercial permit holders. The directed billfish fishery is a recreational fishery. The regulations before this FMP required that all pelagic longline fishermen release any billfish. The regulations were silent on the retention of billfish by other commercial fishermen. NMFS is preferring an alternative that would clarify the regulations and would allow only recreational and charter/headboat fishermen to retain Alantic billfish. General category permit holders participating in a registered HMS tournament could retain billfish during the tournament. Charter/headboat fishermen who

also hold commercial permits (*e.g.*, shark limited access permit) could retain billfish on non-for hire fishing trips only if no HMS on board exceed the recreational limits.

The seventh issue pertains to BFT dealer reports. The preferred alternative would provide an option for BFT dealers to submit certain reports electronically over the Internet once such a system is developed, but would not require it. Although unquantifiable, this alternative is expected to produce positive social and economic impacts for both industry and government, as a result of timesavings incurred when such a system is developed.

The eighth and ninth issues are related to reporting. The preferred alternatives would require no fishing reports and cost-earning reports to be submitted within a certain timeframe and would require either vessel owners or their designee, rather than anglers, to report all non-tournament recreational landings of Atlantic billfish and North Atlantic swordfish. None of these alternatives are expected to have adverse social or economic impacts. Rather, they clarify the regulations and improve data collection.

The tenth issue addresses the Northeast Distant (NED) BFT set-aside for pelagic longline fishermen. NMFS is preferring the alternative that would conduct additional discussions at ICCAT regarding the long-term implications of allowing unused BFT quota from the previous year being added to the subsequent year's allocation. Depending on the results of these discussions the regulations and operation procedures may need to be further amended in the future. In the interim, NMFS would maintain the current regulatory text, but would amend the practice of allowing under/overharvest of this set-aside allocation to be rolled into, or deducted from, the subsequent fishing year's set-aside allocation. This alternative would allow the pelagic longline fishery to retain incidentally caught BFT in the NED to the amount of 25 mt (ww) before landings are counted against the overall Longline category quota. At the proposed stage, NMFS preferred the alternative that would amend the current regulatory text and allow unharvested set-aside quota to be carried forward to subsequent years. That alternative is no longer preferred due to concerns about stockpiling quota and creating potential incentives to target BFT.

The last issue addressed in this section pertains to the inconsistencies between state and Federal regulations. Under the regulations, commercial swordfish and shark fishermen, as a condition of their permit, must abide by Federal regulations when fishing in state waters unless the state has more restrictive regulations. NMFS is preferring an alternative that would expand this permit condition to recreational and charter/headboat fishermen. This alternative is expected to achieve increased consistency between state and Federal regulations for Federally-permitted HMS recreational fishermen, and result in less confusion on behalf of fishermen and improved compliance. Compared with the No Action alternative, the preferred alternative would produce greater ecological benefits with few adverse social and economic impacts.

Essential Fish Habitat

In addition, this Consolidated HMS FMP continues a five-year review of EFH consistent with the EFH guidelines. The Magnuson-Stevens Act requires the Secretary, through NMFS, to establish guidelines to assist in the description and identification of EFH in FMPs, among other

things. The Agency set forth a schedule for the review and update of such EFH identifications based on new scientific evidence or other relevant information. The EFH guidelines articulate processes for determining the extent of EFH for each species and life-stage in a managed fishery. In addition, the EFH guidelines call for periodic review and revision of EFH identified areas based on available information, as well as a complete review of all EFH information at least once every five years. NMFS originally described and identified EFH for all HMS in 1999, and recently updated the EFH for five shark species (blacktip, dusky, finetooth, nurse, and sandbar) in Amendment 1 to the FMP for Atlantic Tunas, Swordfish, and Sharks, which was finalized in 2003. In this document, NMFS includes the information available for all HMS in order to aid in the determination of which species need updates to their EFH identifications. Any updates or resulting changes in management will be done in a future document.

Future Considerations

Beyond the issues addressed in this document or raised during scoping, other new and unresolved matters have been identified by the general public, the HMS and Billfish Advisory Panels, and NMFS staff as important to rebuilding and maintaining fisheries that are economically and biologically sustainable. NMFS may consider these issues or others in future rulemakings. It is important to note that some of these additional issues are complicated, may require specific comments from the public for development (*e.g.*, scoping meetings and/or developmental workshops), and may take several years to complete. These issues include: the BFT fishery (status of BFT, protection of spawning grounds, potential impact of herring fisheries, size limits, filleting at sea); the swordfish fishery (quota underharvests, reporting by recreational anglers, limited access restrictions, time/area closures); the billfish fishery (ESA status review in 2007, stock status, reduction in bycatch and post-release mortality); the shark fishery (new stock assessments, changes to trip limits, limited access restrictions, time/area closures); HMS permit reform; and recordkeeping, reporting, and monitoring of all HMS fisheries. These issues are described in more detail in Section 1.5.

MASTER TABLE OF CONTENTS

Executive Su	mmary	ii
Master Table	e of Contents	xix
Master List o	of Tables	XXX
Master List o	of Figures	li
	nonly Used Abbreviations and Acronyms	
	able of Contents	
-	st of Tables	
-	ion	
	ef Management History	
	ed for Action	
	ectives	
J	nbining Management for Atlantic HMS	
1.4 Con	Implications for Management Measures	
1.4.1	Implications for the Exemption to the Billfish No Sale Provision	
1.4.3	Implications for Highly Migratory Species and Billfish Advisory Panels	
1.4.4	Implications for the FMP Objectives	
	es for Future Consideration and Outlook	
	able of Contents	
-	st of Tables	
-		
-	st of Tables	
-	st of Figures	
·	of Alternatives	
•	atch Reduction	
2.1.1	Workshops	
2.1.1.1		
•	ne, Bottom Longline, and Gillnet Fishermen	
2.1.1.2 2.1.2	HMS Identification Workshops Time/Area Closures	
	ouilding and Preventing Overfishing	
2.2 Red 2.2.1	Northern Albacore Tuna	
2.2.2	Finetooth Sharks	
2.2.3	Atlantic Billfish	
	nagement Program Structure	
2.3.1	Atlantic Bluefin Tuna Quota Management	
2.3.1.1		
2.3.1.2		
2.3.1.3	Inseason Actions	2-45
2.3.2	Timeframe for Annual Management of HMS Fisheries	
2.3.3	Authorized Fishing Gear	
2.3.4	Regulatory Housekeeping	
2.3.4.1		
2.3.4.2		
Chapter 2 Re	eferences	2-83

Chapter 3 Ta	ble of Contents	3-i
Chapter 3 Lis	et of Tables	3-v
_	st of Figures	
-	on of Affected Environment	
	oduction to HMS Management and HMS Fisheries	
3.1.1	History of Atlantic Tunas, Swordfish, and Shark Management	
3.1.1.1	Pre-1999 Atlantic Tunas Management	
3.1.1.2	Pre-1999 Atlantic Swordfish Fishery and Management	
3.1.1.3	Pre-1999 Atlantic Shark Fisheries and Management	
3.1.1.4	1999 Fishery Management Plan for Atlantic Tunas, Swordfish, & Sharks	
3.1.1.5	Post 1999 FMP	
3.1.1.6	Regulatory Amendments Relating to the Pelagic Longline Fishery	
3.1.1.7	Amendment 1 to the Fishery Management Plan for Atlantic Tunas, Swordfish,	
Sharks	3-18	
3.1.1.8	Other Post-1999 FMP Regulations for Atlantic Tunas, Swordfish, and Sharks	3-19
3.1.2	History of Atlantic Billfish Fishery Management	3-21
3.1.2.1	Preliminary Fishery Management Plan (PMP) for Atlantic Billfish and Sharks	
3.1.2.2	The Fishery Management Plan for the Atlantic Billfishes	
3.1.2.3	Interim Rules	
3.1.2.4	Amendment One to the Atlantic Billfish Fishery Management Plan	
3.1.2.5	ICCAT 2000	
3.1.2.6	White Marlin Endangered Species Act (ESA) Listing Review	
3.1.2.7	ICCAT 2002	
3.1.2.8	Recreational Permitting and Reporting Rules	
3.1.2.9		
3.1.2.10		
3.1.3 3.1.4	Summary and Update of Management Measures Taken in 2005 and Early 2006	
	2005 Accomplishments of the International Commission for the Conservation of CAT)	
3.1.4.1	Atlantic Tunas	
3.1.4.1	Atlantic Sharks	
3.1.4.3	Trade and Trade Monitoring.	
3.1.4.4	Data Compliance	
3.1.4.5	Circle Hooks	
	Existing State Regulations	
	us of the Stocks	
3.2.1	Atlantic Swordfish	
3.2.1.1	Life History and Species Biology	
3.2.1.2	Stock Status and Outlook	
3.2.1.3	Effect of Regulations	3-49
3.2.1.4	Recent and Ongoing Research	3-51
3.2.2	Atlantic Bluefin Tuna	3-51
3.2.2.1	Life History and Species Biology	
3.2.2.2	Stock Status and Outlook	
3.2.2.3	Effects of Regulations	
3.2.2.4	Recent and Ongoing Research	
3.2.3	Atlantic BAYS Tuna	
3.2.3.1	Atlantic Bigeye Tuna	
3.2.3.2	Atlantic Yellowfin Tuna	
3.2.3.3	Atlantic Albacore Tuna	3-14

3.2.3.4	Atlantic Skipjack Tuna	3-79
3.2.4	Atlantic Billfish	3-83
3.2.4.1	Blue Marlin	3-83
3.2.4.2	White Marlin	3-92
3.2.4.3	Sailfish	3-98
3.2.4.4	Longbill Spearfish	3-105
3.2.5	Atlantic Sharks	3-107
3.2.5.1	Life History/Species Biology	3-107
3.2.5.2	Stock Status and Outlook	
3.2.5.3	Large Coastal Sharks	3-109
3.2.5.4	Small Coastal Sharks	
3.2.5.5	Pelagic Sharks	3-117
3.2.5.6	Effects of Regulations	3-118
3.2.5.7	Recent and Ongoing Research	3-120
3.3 Hab	itat	3-128
3.3.1	Regulatory Requirements	
3.3.1.1	Habitat Areas of Particular Concern	
3.3.2	Habitat Types and Distributions	
3.3.2.1	Atlantic Ocean	
3.3.2.2	Gulf of Mexico	
3.3.2.3	U.S. Caribbean	
	ery Data Update	
3.4.1	Pelagic Longline Fishery	
3.4.1.1	Domestic History and Current Management	
3.4.1.2	Recent Catch and Landings	
3.4.1.3	Safety Issues	
3.4.1.4	International Issues and Catch	
3.4.2	Purse Seine	
3.4.2.1	Domestic History and Current Management	
3.4.2.2	Recent Catch and Landings	
3.4.2.3	Safety Issues	
3.4.2.4	International Issues and Catch	
3.4.3	Commercial Handgear	
3.4.3.1	Domestic History and Current Management	
3.4.3.2	Recent Catch and Landings	
3.4.3.3	Safety Issues Associated with the Fishery	
3.4.3.4	U.S. vs. International Issues and Catch	
3.4.4	Recreational Handgear	
3.4.4.1	Overview of History and Current Management	
3.4.4.2	Most Recent Catch and Landings Data	
3.4.4.3	Bycatch Issues and Data Associated with the Fishery	
3.4.4.4	Safety Issues Associated with the Fishery	
3.4.4.5	U.S. vs. International Catch	
3.4.5	Bottom Longline	
3.4.5.1	Domestic History and Current Management	
3.4.5.2	Recent Catch and Landings Data	
3.4.5.3	Bottom Longline Bycatch	
3.4.6	Gillnet Fishery	
3.4.6.1	Domestic History and Current Management	
3.4.6.2	Recent Catch and Landings	
3.4.7	Fishery Data: Landings by Species	
~ ,		

3.5 Eco	nomic Status of HMS Fisheries	3-228
3.5.1	Commercial Fisheries	3-229
3.5.1.1	Ex-Vessel Prices	3-230
3.5.1.2	Revenues	3-237
3.5.1.3	Wholesale Market	3-239
3.5.2	Recreational Fisheries	
3.6 Con	nmunity and Social Update	3-244
3.6.1	Overview of Current Information and Rationale	
3.6.2	Social Impacts of Selected 2005 Regulatory Actions	
3.6.3	Summary of New Social and Economic Data Available	
3.6.3.1	2005 Social Science Publications	
3.6.3.2	Summary of Social Data and Information for FEIS	
3.6.3.3	HMS Community Profile Needs	
3.7 Inter	rnational Trade and Fish Processing	
3.7.1	Overview of International Trade for Atlantic HMS	
3.7.1.1	Trade Monitoring	
3.7.1.2	Bluefin Tuna Statistical Document	
3.7.1.3	Swordfish Certificate of Eligibility and Statistical Document	
3.7.1.4	Bigeye Tuna Statistical Document	
3.7.1.5	Yellowfin Tuna Form 370	
3.7.1.6	Billfish Certificate of Eligibility	
3.7.2	U.S. Exports of HMS	
3.7.2.1	Atlantic and Pacific Bluefin Tuna Exports	
3.7.2.2	Other Tuna Exports	
3.7.2.3	Shark Exports	
3.7.2.4	Re-exports of Atlantic HMS	
3.7.2.5	Summary of Atlantic HMS Exports	
3.7.3	U.S. Imports of Atlantic HMS	
3.7.3.1	Bluefin Tuna Imports	
3.7.3.2	Other Tuna Imports	
3.7.3.3	Swordfish Imports	
3.7.3.4	Shark Imports	
3.7.3.5	Summary of U.S. Imports of Atlantic HMS	
3.7.4	The Use of Trade Data for Conservation Purposes	
3.7.5	Overview of the Processing Industry for Atlantic HMS	
3.7.5.1	Processing and Wholesale Sectors	
3.8 Byc	atch, Incidental Catch, and Protected Species	
3.8.1	Bycatch Reduction and the Magnuson-Stevens Act	
3.8.2	Standardized Reporting of Bycatch	
3.8.2.1	U.S. Atlantic Pelagic Longline Fishery	
3.8.2.2	Purse Seine Fishery	
3.8.2.3	Shark Bottom Longline Fishery	
3.8.2.4	Shark Gillnet Fishery	
3.8.2.5	Commercial Handgear Fishery	
3.8.2.6	Recreational Handgear Fishery	
3.8.3	Bycatch Reduction in HMS Fisheries	
	S Permits and Tournaments	
3.9.1	Upgrading and Safety Issues	
3.9.2	Atlantic Tunas Permits.	
3.9.3	HMS CHB Permits	

3.9.6	HMS Angling Permit	3-278
3.9.4	Dealer Permits	3-278
3.9.5	Exempted Fishing Permits (EFPs), Display Permits, Chartering Permits,	and Scientific
Research	n Permits (SRPs)	
3.9.6	Atlantic HMS Tournaments	3-283
3.9.7	Evaluation and Monitoring of Bycatch	3-291
3.9.8	Bycatch Mortality	3-291
3.9.8.	1 Introduction	3-291
3.9.8.	2 Mortality by Fishery	3-292
3.9.8.	3 Code of Angling Ethics	3-294
3.9.9	Interactions of HMS Fishing Gears with Protected Species	3-294
3.9.9.	1 Interactions and the Marine Mammal Protection Act	3-295
3.9.9.	2 Interactions and the ESA	3-297
3.9.9.	3 Interactions with Seabirds	3-300
3.9.10	Measures to Address Protected Species Concerns	
3.9.11	Bycatch of HMS in Other Fisheries	3-301
3.9.11	.1 Squid Mid-Water Trawl	3-302
3.9.11	.2 Menhaden Purse Seine Fishery	3-302
3.9.11	.3 Shrimp Trawl Fishery	3-303
3.9.11	.4 Southeast Gillnet Fishery	3-304
3.9.12	Effectiveness of Existing Time/Area Closures in Reducing Bycatch	3-304
3.9.12	Prohibition of Live Bait in the Gulf of Mexico	3-305
3.9.12	2.2 Conclusions	3-306
3.9.13	Evaluation of Other Bycatch Reduction Measures	3-306
Chapter 3 R	eferences	3-309
-	able Of Contents	
-	ist of Tables	
-	ist of Figures	
.0 Environ	mental Consequences of Alternatives	4-1
	catch Reduction	
4.1.1		
4.1.1.	•	
	ine, Bottom Longline, and Gillnet Fishermen	
	2 HMS Identification Workshops	
4.1.2	Time Area Closures	
	building and Preventing Overfishing	
4.2.1	Northern Albacore Tuna	
4.2.2	Finetooth Sharks	
4.2.3	Atlantic Billfish	
	inagement Program Structure	
4.3.1	Atlantic Bluefin Tuna Quota Management	
4.3.1.	· · · · · · · · · · · · · · · · · · ·	
4.3.1.		
4.3.1.		
4.3.1.	Timeframe for Annual Management of HMS Fisheries	
4.3.3	Authorized Fishing Gear	
4.3.4	1 1641011204 1 19111115 0041	437
	Regulatory Housekeeping	4-254
44 Im	Regulatory Housekeeping	
4.4 Im 4.4.1	Regulatory Housekeepingpacts on Essential Fish Habitat	4-289

4.4.2		
4.4.3	Northern Atlantic Albacore Tuna	4-290
4.4.4	Finetooth Sharks	4-290
4.4.5	· · · · · · · · · · · · · · · · · · ·	
4.4.6	$\boldsymbol{\mathcal{E}}$	
4.4.7		
4.4.8	ϵ	
4.4.9		
4.5	Impacts on Protected Resources	
4.6	Environmental Justice	
4.7	Coastal Zone Management Act	
4.8	Cumulative Impacts	
4.8.1	····, ····, ···· · · · · · · · · · · ·	
4.8.2	F	
4.8.3	Cumulative Economic and Social Impacts	4-304
Chapter	4 References	4-323
Chapter	5 Table of Contents	5-i
-	ation and Unavoidable Impacts	
5.0 minug 5.1	Mitigation Measures	
5.2	Unavoidable Adverse Impacts	
5.3	Irreversible and Irretrievable Commitment of Resources	
_	6 Table of Contents	
-	6 List of Tables	
6.0 Econ	omic Evaluation	6-1
6.1	Number of Fishing and Dealer Permit Holders	
6.1.1	6	
6.1.2		
6.1.3		
6.2	Gross Revenues of Fishermen	
6.3	Variable Costs and Net Revenues	
6.4	Expected Economic Impacts of the Alternatives	6-4
	Bycatch Reduction	
	4.1.1 Workshops	
	4.1.2 Time/Area Closures	
6.4.2	8	
	4.2.1 Northern Albacore	
	4.2.2 Finetooth Sharks	
	4.2.3 Atlantic Billfish	
6.4.3		
	4.3.1 Bluefin Tuna Quota Management	
	4.3.2 Timeframe for Annual Management of HMS Fisheries	
	4.3.4 Regulatory Housekeeping	
-	6 References	
-	7 Table of Contents	
Chapter	7 List of Tables	7-i
7.0 Regu	latory Impact Review	7-1

7.1	Description of the Management Objectives	7-1
7.2	Description of the Fishery	
7.3	Statement of the Problem	7-1
7.4	Description of Each Alternative	7-1
7.5	Economic Analysis of Expected Effects of Each Alternative Relative to the	e Baseline 7-
7.6	Conclusion	7-3
Chapter	8 Table of Contents	
-	Regulatory Flexibility Analysis	
8.1	Statement of the Need for and Objectives of this Final Rule	
8.2	A Summary of the Significant Issues Raised By the Public Comments in R	
0.2	the Initial Regulatory Flexibility Analysis, a Summary of the Assessmen	
	Agency of Such Issues, and a Statement of Any Changes Made in the R	
	Result of Such Comments	
8.2.1		
8.2.2	<u>*</u>	
8.2.3		
8.2.4	Atlantic Billfish	8-4
8.2.5	Atlantic Bluefin Tuna Domestic Management Measures	8-6
8.2.6	\mathcal{E}	
8.2.7	\mathcal{E}	
8.2.8		
8.3	Description and Estimate of the Number of Small Entities to Which the Pr	-
	Will Apply	
8.3.1	- T	
8.3.2		
8.3.3		
8.3.4		
8.3.5 8.3.6		
8.3.7	<u>e</u>	
8.3.8	C	
8.3.9	ϵ	
	Description of the Projected Reporting, Record-Keeping, and Other Comp	
0	Requirements of the Proposed Rule, Including an Estimate of the Classe	
	Entities Which Will Be Subject to the Requirements of the Report or Re	
8.5	Description of the Steps the Agency Has Taken to Minimize the Significant	
0.5	Impact on Small Entities Consistent with the Stated Objectives of Appli	
	Statutes, Including a Statement of the Factual, Policy, and Legal Reason	
	Selecting the Alternative Adopted in the Final Rule and the Reason Tha	
	of the Other Significant Alternatives to the Rule Considered by the Age	
	Affect Small Entities Was Rejected	•
8.5.1	<u> </u>	
	5.1.1 Workshops	
	5.1.2 Time/Area Closures	
8.5.2		
	5.2.1 Northern Albacore Tuna	
	5.2.2 Finetooth Sharks	

8.5.2.3	3 Atlantic Billfish	8-16
8.5.3	Management Program Structure	8-18
8.5.3.	1 Bluefin Tuna Quota Management	8-18
8.5.3.2	\mathcal{C}	
8.5.3	3 Authorized Fishing Gears	8-19
8.5.3.4	4 Regulatory Housekeeping	8-20
Chapter 9 T	able of Contents	9-i
-	ist of Tables	
_	ist of Figures	
-	nity Profiles	
	roduction	
	ethodology	
9.2.1	Previous community profiles and assessments	
9.2.2	Information Used in this Assessment	
9.2.2.		
	ited States Demographic Profile	
	ate and Community Profiles	
9.4.1	Maine	
9.4.2	New Hampshire	
9.4.3	Massachusetts	
9.4.3.		
9.4.3.2		
9.4.4	Rhode Island	
9.4.5	Connecticut	
9.4.6	New York	
9.4.7	New Jersey	
9.4.7.	· · · · · · · · · · · · · · · · · · ·	
9.4.7.2		
9.4.8	Delaware	
9.4.9	Maryland	9-29
9.4.10	Virginia	
9.4.11	North Carolina	9-31
9.4.11	.1 Hatteras	9-33
9.4.11		
9.4.12	South Carolina	
9.4.13	Georgia	
9.4.14	Florida	
9.4.14	1 '	
9.4.14		
9.4.14	,	
9.4.14	3 /	
9.4.14	,	
9.4.15	Alabama	
9.4.16	Mississippi	
9.4.17	Louisiana	
9.4.17	· · · · · · · · · · · · · · · · · · ·	
9.4.17		
9.4.18	Texas	
9419	Puerto Rico	9-65

9.	4.19.1 Arecibo, Puerto Rico	9-66
9.5	Future Assessments	9-68
Chapter	9 References	9-76
-	10 Table of Contents	
_	10 List of Tables	
-	10 List of Tables	
_	ssential Fish Habitat	
10.0 E		
	EFH Five-Year Review Process	
10.2	1	
10.2	1	
10.3 10.3	Summary of Review and Findings 1 Tunas	
10.3		
10.3		
10.3		
	Threats to Essential Fish Habitat	
10.4		
10.4		
	0.4.2.1 HMS EFH Overview	
	0.4.2.2 Impact of HMS Federally regulated gear on HMS and non-HMS EFH	
10.4		
10.4		
10.4	.5 Non-Federally Managed Fishing Activities	10-29
10.4		
10.4	.7 Summary	10-45
10.5	6 I	
10	0.5.1.1 Land-based Activities That May Impact HMS EFH	
10	0.5.1.2 Coastal and Offshore Activities That May Impact HMS EFH	
	.2 Cumulative Impacts	
Chapter	10 References	10-54
Chapter	s 11-13 Table of Contents	11-i
_	Other Considerations	
11.1	National Standards	
11.2	Ongoing Management and the Procedure for Adjusting Management Measures.	
11.3	Consideration of Magnuson-Stevens Act Section 304(g) Measures	
	ist of Preparers	
	•	
	ist Of Agencies, Organizations, And Persons Consulted And To Whom Copi	
	he EIS Will Be Sent	
Appendi	x A Table of Contents	A-i
Appendi	x A List of Tables	A-i
Appendi	x A List of Figures	A-vi
	endix: Time/Area Closures: additional analyses and results	
	x B Table of Contents	
	x B List of Tables	
Appendi	x B List of Figures	B-iii

B Appendix: Essential Fish Habitat	B-1
B.1 Life History Accounts and Essential Fish Habitat Descriptions	
B.1.1 Tuna	
B.1.1.1 Atlantic Albacore Tuna	
B.1.1.2 Atlantic Bigeye Tuna	
B.1.1.3 Atlantic Bluefin Tuna	
B.1.1.4 Atlantic Skipjack Tuna	
B.1.1.5 Atlantic Yellowfin Tuna	
B.1.2 Swordfish	
B.1.3 Billfish	
B.1.3.1 Blue Marlin	
B.1.3.2 White Marlin	B-15
B.1.3.3 Sailfish	B-17
B.1.3.4 Longbill Spearfish	
B.1.4 Large Coastal Sharks	
B.1.4.1 Basking Sharks	B-20
B.1.4.2 Hammerhead Sharks	B-21
B.1.4.3 Mackerel Sharks	B-23
B.1.4.4 Nurse Sharks	B-24
B.1.4.5 Requiem Sharks	B-26
B.1.4.6 Sand Tiger Sharks	B-39
B.1.4.7 Whale Sharks	B-40
B.1.4.8 Small Coastal Shark	B-41
B.1.4.9 Hammerhead Sharks	B-42
B.1.4.10 Requiem Sharks	
B.1.5 Pelagic Sharks	
B.1.5.1 Cow sharks	
B.1.5.2 Mackerel Shark	
B.1.5.3 Requiem Sharks	
B.1.5.4 Thresher Sharks	B-54
Appendix B References	B-196
Appendix C List of Tables	
C Appendix: Aggregate Domestic Pelagic Longline and Recreational	
Marlin Fishing Mortality Estimates 2001 - 2004	
Appendix D: Table of Contents	
D Proposed Rule and DEIS Comments and Responses	
D.1 Bycatch Reduction	
D.1.1 Workshops	
D.1.2 Time/Area Closures	
D.2 Rebuilding and Preventing Overfishing	D-46
D.2.1 Northern Albacore Tuna	
D.2.2 Finetooth Sharks	
D.2.3 Atlantic Billfish	D-58
Gears and Gear Restrictions	
Circle Hooks And/Or Post-Release Mortality Data	
Elimination of the 'No Sale' Exemption	
General Billfish Comments	
D.3 Management Program Structure	
D.3.1 Bluefin Tuna Quota Management Measures	

D.3	3.2	Timeframe for Annual Management of HMS Fisheries	D-103
D.3	3.3	Authorized Fishing Gear	D-105
D.3	3.4	Regulatory Housekeeping	D-127
D.4	Esse	ential Fish Habitat	D-145
D.5	Eco	nomic and Social Impacts	D-151
D.6		solidation of the FMPs	
D.7	Obj	ectives of the FMP	D-156
D.8		nment Period/Outreach	
D.9	Gen	eral	D-160
D.9		Recreational	
D.9	9.2	Commercial Fishery	D-165
D.9	9.3	Longline	D-166
D.9	9.4	Swordfish	D-167
D.9	9.5	Tunas	D-170
D.9	9.6	Sharks	D-170
D.9	9.7	Fishing Mortality and Bycatch Reduction	D-174
D.9	9.8	Permitting, Reporting, and Monitoring	D-180
D.9	9.9	Enforcement	D-183
D.9	9.10	ICCAT	D-185
APPEN	DIX 1	E TABLE OF CONTENTS	E-i
E OM	B PE	ER REVIEW BULLETIN	E-1
E.1	Pee	Review by Gregory Skomal, Commonwealth of Massachusetts, Divi	sion of
		farine Fisheries, December 21, 2005	
E.2		ponse to OMB Peer Review by Gregory Skomal, Commonwealth of	
		Assachusetts, Division of Marine Fisheries, December 21, 2005	E-15
E.3		Review by Chris Boggs and Keith Bigelow, NMFS SWFSC, January	
E.4		ponse to OMB Peer Review by Chris Boggs and Keith Bigelow, NMF	
Д. 1	-	anuary 9, 2006	
			L-23
F 5		Review by Paul I Rago NMES NEESC January 25, 2006	E 33
E.5	Peer	Review by Paul J. Rago, NMFS NEFSC, January 25, 2006	
E.5 E.6	Peer	ponse to OMB Peer Review by Paul J. Rago, NMFS NEFSC, January	

MASTER LIST OF TABLES

Table 1	The preferred alternatives at the draft and final stage of the Consolidated HMS FMP and the expected implementation date
Table 1.1	Table indicating whether actions in this document are amending the FMP or are being taken as framework actions
Table 1.2	Current Advisory Panel Seat Allocation1-13
Table 1.3	Previous and Final Objectives of the Atlantic Tunas, Swordfish, and Shark, Billfish, and Consolidated HMS FMPs. <i>Italicized</i> text indicates the differences in objectives between the two previous FMPs
Table 2.1	Summary and status of activities, anticipated results, and associated timelines for preventing overfishing of finetooth sharks
Table 2.2	List of Proposed Regulatory Changes2-59
Table 3.1	Summary of 2005 Federal Register Notices Related to HMS3-28
Table 3.2	State Rules and Regulations Pertaining to Atlantic HMS, as of May 30, 2006.3-33
Table 3.3	Stock Assessment Summary Table. Source: SCRS, 2004 and 2005, Cortes, 2002, and Cortes <i>et al.</i> 2002
Table 3.4	Atlantic Swordfish Stock Summary (weights given in mt ww). Source: SCRS, 2005
Table 3.5	Summary Table for the Status of West Atlantic Bluefin Tuna. Source: ICCAT, 2005
Table 3.6	Summary Table for the Status of East Atlantic Bluefin Tuna. Source: ICCAT, 2005
Table 3.7	Probability of western Atlantic bluefin tuna achieving rebuilding target by 2018. Source: ICCAT, 2004
Table 3.8	Summary Table for the Status of Atlantic Bigeye Tuna. Source: ICCAT, 20053-66
Table 3.9	Summary Table for the Status of Atlantic Yellowfin Tuna. Source: ICCAT, 2005.
Table 3.10	Summary Table for the Status of North Atlantic Albacore Tuna. Source: ICCAT, 2005
Table 3.11	Summary Table for the Status of South Atlantic Albacore Tuna. Source: ICCAT, 2005
Table 3.12	Summary Table for the Status of West Atlantic Skipjack Tuna. Source: ICCAT, 2005
Table 3.13	Summary of Atlantic Blue Marlin Stock Assessment data. Weights are in metric tons, whole weight. Source: SCRS, 2005
Table 3.14	Summary of Atlantic White Marlin Stock Assessment data. Weights are in metric tons, whole weight. Source: SCRS, 2005
Table 3.15	Summary of Atlantic Sailfish Stock Assessment data. Weights are in metric tons, whole weight. Source: SCRS, 2004
Table 3.16	Common names of shark species included within the four species management units under the purview of the HMS management division

Table 3.17	Summary Table of Biomass and Fishing Mortality for Large Coastal Sharks (LCS). Source: Cortes <i>et al.</i> , 2002
Table 3.18	Summary Table of Biomass and Fishing Mortality for Small Coastal Sharks (SCS) Source: Cortes, 2002
Table 3.19	Summary table of the status of the biomass of finetooth sharks. Sources: 2002 SCS stock assessment; E. Cortes, personal communication. LRSG=lagged recruitment, survival, and growth model; SPM=surplus production model3-116
Table 3.20	Summary table of the status of the biomass of finetooth sharks. Sources: 2002 SCS stock assessment; E. Cortes, personal communication. LRSG=lagged recruitment, survival, and growth; SPM=surplus production model3-116
Table 3.21	Calendar Year 2004 U.S. vs International Catch of HMS (mt ww) other than sharks. Source: SCRS, 2005.
Table 3.22	Average Number of Hooks per Pelagic Longline Set, 1999-2004. Source: Data reported in pelagic longline logbook
Table 3.23	Observer Coverage of the Pelagic Longline Fishery. Source: Yeung, 2001; Garrison, 2003; Garrison and Richards, 2004; Garrison, 2005; Walsh and Garrison, 2006
Table 3.24	Reported Catch of Species Caught by U.S. Atlantic Pelagic Longlines, in Number of Fish, for 1999-2004. Source: Pelagic Longline Logbook Data3-150
Table 3.25	Reported Landings in the U.S. Atlantic Pelagic Longline Fishery (in mt ww) for 1999-2004. Source: NMFS, 2004a; NMFS, 2005
Table 3.26	Summary of Marine Mammal Interactions in the Pelagic Longline Fishery, 1992-1998. Source: Yeung, 1999a; Yeung, 1999b
Table 3.27	Summary of Marine Mammal Interactions in the Pelagic Longline Fishery, 1999-2005. Sources: Yeung, 2001; Garrison, 2003; Garrison and Richards, 2004; Garrison, 2005; Walsh and Garrison, 2006
Table 3.28	Estimated number of leatherback and loggerhead sea turtle interactions in the U.S. Atlantic pelagic longline fishery, 2001-2005 by statistical area. Sources: Walsh and Garrison, 2006; Garrison, 2005; Garrison and Richards, 2004; Garrison 2003.
Table 3.29	Seabird Bycatch in the U.S. Atlantic Pelagic Longline Fishery, 1992-2005. Source: NMFS, 2004a; NMFS PLL fishery observer program (POP) data 3-158
Table 3.30	Status of Seabird Bycatch in the U.S. Atlantic Pelagic Longline Fishery, 1992-2005. Source: NMFS PLL fishery observer program (POP) data3-159
Table 3.31	Preliminary Expanded Estimates of Seabird Bycatch (D = discarded dead and A = discarded alive) and bycatch rates (all seabirds per 1,000 hooks) in the U.S. Atlantic pelagic longline fishery, 1997-2004. Source: NMFS, 2004a; NMFS PLL fishery observer program (POP) data
Table 3.32	ICCAT Bycatch Table (LL, longline; GILL, gillnets; PS, purse-seine; BB, baitboat; HARP, harpoon; Trap, traps). Source: SCRS, 20043-165
Table 3.33	Estimated International Longline Landings of HMS, Other than Sharks, for All Countries in the Atlantic: 1999-2004 (mt ww) ¹ . Source: SCRS, 20053-166

Table 3.34	Nominal Catches of Blue Shark Reported to ICCAT (landings and discards in t) by Major Gear and Flag between 1990 and 2002. Source: SCRS, 2004; SCRS, 2005
Table 3.35	Nominal Catches of Shortfin Mako Shark Reported to ICCAT (landings and discards in t) by Major Gear and Flag between 1990 and 2002. Source: SCRS, 2004; SCRS, 2005
Table 3.36	Domestic Atlantic Tuna Landings for the Purse Seine Fishery: 1999-2004 (mt ww). Northwest Atlantic Fishing Area. Source: U.S. National Report to ICCAT: 2005
Table 3.37	Estimated International Purse Seine Atlantic Tuna Landings in the Atlantic and Mediterranean: 1999-2004 (mt ww). Source: SCRS, 2005
Table 3.38	Domestic Landings for the Commercial Handgear Fishery, by Species and Gear, for 1999-2004 (mt ww). Source: U.S. National Report to ICCAT: 2005 3-177
Table 3.39	Domestic Landings for the Commercial Handgear Fishery by Species and Region for 1999-2004 (mt ww). Source: U.S. National Report to ICCAT: 2005 3-178
Table 3.40	Estimated number of vessel trips targeting large pelagic species, 2001-2004. Source: Large Pelagics Survey database
Table 3.41	Updated Domestic Landings for the Atlantic Tunas, Swordfish and Billfish Recreational Rod and Reel Fishery, 1997-2004 (mt ww)*. Sources: NMFS, 2004; NMFS, 2005. (Recreational shark landings are provided in Table 3.44 through Table 3.47)
Table 3.42	Preliminary RBS Recreational Billfish Landings in numbers of fish (calendar year). Source: NMFS Recreational Billfish Survey (RBS)
Table 3.43	Number of billfish reported to NMFS via call-in system by fishing year, 2002-2005. Source: G. Fairclough, pers. comm
Table 3.44	Estimates of Total Recreational Harvest of Atlantic Sharks: 1998-2004 (numbers of fish in thousands). Source: 1998-2000 (Cortés, pers. comm.); 2001-2004 (Cortés, 2005a; 2005b). Estimates for 2001-2004 do not include prohibited species
Table 3.45	Recreational Harvest of Atlantic Large Coastal Sharks (LCS) by Species, in number of fish: 1998-2004. Sources: 1998-2000 (Cortés, pers. comm.); 2001-2004 (Cortés, 2005a; 2005b). Total estimates for 2001-2004 do not include prohibited species
Table 3.46	Recreational Harvest of Atlantic Pelagic Sharks by Species, in number of fish: 1998-2004. Sources: 1998-2000 (Cortés, pers. comm.); 2001-2004 (Cortés, 2005a; 2005b). Total estimates for 2001-2004 do not include prohibited species.3-189
Table 3.47	Recreational Harvest of Atlantic SCS by Species, in number of fish: 1998-2004. Source: 1998-2000 (Cortés, pers. comm.); 2001-2004 (Cortés, 2005a; 2005b). Total estimates for 2001-2004 do not include prohibited species
Table 3.48	Observed or reported number of HMS kept ¹ and released in the rod and reel fishery, Maine through Virginia, 1997-2004. Source: Large Pelagic Survey (LPS) Preliminary Data

Table 3.49	Total 2004 Reported Boating Accident Types. Source: USCG Boating Statistics, 2004
Table 3.50	Overall 2004 Reported Boating Accident Cause-of-Death Statistics. Source: USCG Boating Statistics, 2004
Table 3.51	Species composition of observed bottom longline catch during 2003. Source: Burgess and Morgan, 2004
Table 3.52	Species composition of observed bottom longline catch during 2004. Source: Burgess and Morgan, 2005
Table 3.53	Directed bottom longline shark observed catch and disposition, 2003. Source: Burgess and Morgan, 2004
Table 3.54	Total number of Observed Sea Turtle Interactions by Species by Month for Years 1994-2006 in the Shark Bottom Longline Fishery. Source: Shark Bottom Longline Observer Program
Table 3.55	Total number of Observed Sea Turtle Interactions by Year for Years 1994-2006 in the Shark Bottom Longline Fishery. Source: Shark Bottom Longline Observer Program. Letters in parentheses indicate whether the sea turtle was released alive (A), dead (D), or in an unknown (U) condition
Table 3.56	Total Strikenet Shark Catch and Bycatch by Species in order of Decreasing Abundance for all Observed Trips, 2003. Source: Carlson and Baremore, 20033-211
Table 3.57	Total Shark Catch by Species and Species Disposition in Order of Decreasing Abundance for all Observed Driftnet Sets, 2003. Source: Carlson and Baremore, 2003
Table 3.58	Total bycatch in NMFS observed drift gillnet sets in order of decreasing abundance and species disposition for all observed trips, 2003. Source: Carlson, 2003
Table 3.59	Total number of Observed Sea Turtle Interactions by Year from 2000-2006 in the Shark Gillnet Fishery. Source: Directed Shark Gillnet Observer Program. Letters in parentheses indicate whether the sea turtle was released alive (A), dead (D), or unknown (U)
Table 3.60	Protected Species Interactions in Drift Gillnet Sets During the Directed Shark Gillnet Fishery for All Observed Trips, 2003. Source: Carlson, 20033-213
Table 3.61	U.S. Landings (mt) of Bluefin Tuna by Gear and Area, 1997-2004. Source: NMFS, 2005
Table 3.62	U.S. Landings (mt) of Yellowfin Tuna by Gear and Area, 1997-2004. Source: NMFS, 2005
Table 3.63	U.S. Landings (mt) of Skipjack Tuna by Gear and Area, 1997-2004. Source: NMFS, 2005
Table 3.64	U.S. Landings (mt) of Bigeye Tuna by Area and Gear, 1997-2004. Source: NMFS, 2005
Table 3.65	U.S. Landings (mt) of Albacore Tuna by Gear and Area, 1997-2004. Source: NMFS, 2005
Table 3.66	U.S. Catches and Landings (mt) of Swordfish by Gear and Area, 1997-2004. Source: NMFS, 2005

Table 3.67	U.S. Landings (mt) and dead discards of Blue Marlin, White Marlin and Sailfish by Gear and Area, 1998-2002. Source: NMFS, 2003
Table 3.68	Commercial landings of large coastal sharks in lb dw: 1999-2004. Sources: Data from 1999-2001, Cortés pers. Comm.; data from 2002-2003, Cortés 2003; Cortés and Neer, 2005
Table 3.69	Commercial landings of small coastal sharks in lb dw: 1999-2004. Source: Cortés and Neer, 2002; Cortés, 2003. Cortés and Neer, 2005
Table 3.70	Commercial landings of pelagic sharks in lb dw: 1999-2004. Sources: Data from 2000-2001, Cortés pers. comm.; Cortés, 2003; Cortés and Neer, 2005 3-223
Table 3.71	Estimates of total landings and dead discards for large coastal sharks from 1981 through 2004 (numbers of fish in thousands). Modified from the 1998 and 2002 Report of the Shark Evaluation Workshop (NMFS 1998, 2002), Cortés and Neer (2002), and Cortés (2003, 2005)
Table 3.72	Commercial landings of LCS (including unclassified sharks) in the Atlantic and Gulf of Mexico by region and year in mt dw for QMS and Logbook data and mt www for Canvass and CFDBS data from 1999-2003
Table 3.73	Commercial landings of SCS in the Atlantic and Gulf of Mexico by region and year year in mt dw for QMS and Logbook data and mt ww for Canvass and CFDBS data from 1999-2003.
Table 3.74	Inflation Price Indexes. The CPI-U is the standard Consumer Price Index for all urban consumers (1982-1984=100) produced by U.S. Department of Labor Bureau of Labor Statistics. The source of the Producer Price Index (PPI) for unprocessed finfish (1982=100) is also the Bureau of Labor Statistics. The Gross Domestic Product Implicit Price Deflator (200=100) is produced by the U.S. Department of Commerce Bureau of Economic Analysis and obtained from the Federal Reserve Bank of St. Louis (http://www.stlouisfed.org/)
Table 3.75	Average ex-vessel prices per lb dw for Atlantic HMS by gear and area. Source: Dealer weighout slips from the Southeast Fisheries Science Center and Northeast Fisheries Science Center, and bluefin tuna dealer reports from the Northeast Regional Office. HND=Handline, harpoon, spears, trot lines, and trolls, PLL=Pelagic longline, BLL=Bottom longline, Net=Gillnets and pound nets, TWL=Trawls, SEN=Seines, TRP=Pots and traps, DRG=Dredge, and UNK=Unknown. Gulf of Mexico includes: TX, LA, MS, AL, and the west coast of FL. S. Atlantic includes: east coast of FL. GA, SC, and NC dealers reporting to Southeast Fisheries Science Center. Mid-Atlantic includes: NC dealers reporting to Northeast Fisheries Science Center, VA, MD, DE, NJ, NY, and CT. N. Atlantic includes: RI, MA, NH, and ME. For bluefin tuna, all NC landings are included in the Mid-Atlantic
Table 3.76	Average ex-vessel prices per lb for Atlantic HMS by area
Table 3.77	Estimates of the total ex-vessel annual revenues of Atlantic HMS fisheries. Sources: NMFS, 1997; NMFS 2004a; Cortes, 2003; and bluefin tuna dealer reports from the Northeast Regional Office
Table 3.78	The overall average wholesale price per lb of fresh HMS sold in Atlantic and Gulf of Mexico states as reported by the Fulton Fish Market. Source: NMFS, 20043-239

Table 3.79	Average Atlantic HMS charterboat rates for day trips. Source: NMFS searches for advertised daily charter rates of HMS Charter/Headboat permit holders. (Observations=99)
Table 3.80	United States exports of Atlantic and Pacific bluefin tuna, 1999-2004. Sources: NMFS BSD Program, NERO, and Census Bureau
Table 3.81	Amount and value of U.S. exports of albacore tuna from all ocean areas, 1999-2004 (Census Bureau data) and U.S. landings of North Atlantic albacore tuna (2005 U.S. National Report to ICCAT)
Table 3.82	Amount and value of U.S. exports of yellowfin tuna from all ocean areas, 1999-2004 (Census Bureau data) and U.S. landings of Atlantic yellowfin tuna (2005 U.S. National Report to ICCAT)
Table 3.83	Amount and value of U.S. exports of skipjack tuna from all ocean areas, 1999-2004 (Census Bureau data) and U.S. landings of West Atlantic skipjack tuna (2005 U.S. National Report to ICCAT)
Table 3.84	Amount and value of U.S. exports of bigeye tuna from all ocean areas, 1999-2004 (Census Bureau data) and U.S. landings of Atlantic bigeye tuna (2005 U.S. National Report to ICCAT)
Table 3.85	Amount and value of U.S. shark product exports from 1999-2004. Source: Census Bureau
Table 3.86	Imports of Atlantic and Pacific bluefin tuna into the United States: 1999-2004. Sources: NMFS BSD program and CBP data
Table 3.87	Imports of bigeye tuna into the United States from all ocean areas combined: 2001-2004. Source: Census Bureau data
Table 3.88	Imports of yellowfin tuna into the United States from all ocean areas combined: 1999-2004. Source: Census Bureau data
Table 3.89	Imports of albacore tuna into the United States from all ocean areas combined: 1999-2004. Source: Census Bureau data
Table 3.90	Imports of skipjack tuna from all ocean areas combined into the United States: 1999-2004. Source: U.S. Census Bureau data
Table 3.91	Swordfish import data for the 2004 calendar year collected under the NMFS Swordfish Import Monitoring Program
Table 3.92	Imported swordfish products by year: 1999-2004. Source: Census Bureau data3-260
Table 3.93	U.S. imports of shark products from all ocean areas combined: 1999-2004. Source: Census Bureau data
Table 3.94	Summary and current status of ICCAT recommended trade sanctions for bluefin tuna, swordfish, and bigeye tuna implemented by the United States
Table 3.95	Distribution of Shark, Swordfish, and Tuna longline Limited Access Permits Between 2001 and 2006. Data for 2001-2005 are as of October 1 for each year3-275
Table 3.96	The number of Atlantic tuna permit holders in each category as of October 2001 through 2005. Permit numbers for 2006 are as of February 1, 2006. The actual number of 2006 permit holders in each category is subject to change as individuals renew or allow their permits to expire

Table 3.97	CHB Permits by State as of February 1, 2006
Table 3.98	Number of shark and swordfish dealer permits issued in each state or country as of October 2001-2005. Permits for 2006 are as of February 1, 2006. The actual number of permits per may change as permit holders move or sell their businesses
Table 3.99	Number of Atlantic tuna dealer permits by state issued in the 2005 calendar year. Dealers may obtain a permit to sell and purchase only bluefin tuna, only BAYS tunas, or both bluefin and BAYS tunas
Table 3.100	Number of International Trade Permits (ITP) by state (province) as of February 1, 2006.
Table 3.101	Number of Exempted Fishing Permits (EFPs), Display Permits, and Scientific Research Permits (SRPs) issued between 2002 and 2006
Table 3.102	Number of Registered HMS Tournaments by State between 2001 and 2005. Source: NMFS Atlantic HMS Tournament Registration Database
Table 3.103	Number and Percent of All 2005 HMS Tournaments Awarding Points or Prizes for an HMS. Source: NMFS Atlantic HMS Tournament Registration Database3-286
Table 3.104	Registered Blue Marlin Tournaments, 2005. Source: NMFS Atlantic HMS Tournament Registration Database
Table 3.105	Registered White Marlin Tournaments, 2005. Source: NMFS Atlantic HMS Tournament Registration Database
Table 3.106	Registered Sailfish Tournaments, 2005. Source: NMFS Atlantic HMS Tournament Registration Database
Table 3.107	Summary of bycatch species in HMS fisheries, Marine Mammal Protection Act (MMPA) category, endangered Species Act (ESA) requirements, data collection, and management measures by fishery/gear type. (Excerpted from HMS Bycatch Priorities and Implementation Plan and updated through May 2006)
Table 3.108	Atlantic HMS Landed (mt ww) Incidental to Trawl Fisheries, 1998-2004. Source: NMFS, 2003, NMFS, 2005
Table 3.109	Expanded estimates of bycatch (number of fish) of bonnethead, Atlantic sharpnose, and finetooth sharks in the U.S. south Atlantic shrimp trawl fishery based on within stratum expansion by effort as trips by fishing year. Source: Cortés, 2002
Table 3.110	Estimates (in thousands of individuals and pounds dressed weight) of the bycatch of small coastal sharks (as a complex and by species) in the shrimp trawl fishery operating in the Gulf of Mexico. Source: S. Nichols, NMFS Pascagoula Lab., pers. comm. as cited in Cortés, 2002
Table 3.111	Comparison of the number of hooks reported set in the Gulf of Mexico with dead or live bait, or a combination of both baits, 1999-2004 (numbers in parentheses are percent of the total number of hooks set in the Gulf of Mexico). Source: PLL Logbook data
Table 4.1	Extrapolated Total Sea Turtle and Marine Mammal Interactions in the HMS PLL fishery for 1999-2005 and the 3 Year ITSs for 2004-2006 and 2007-2009.

	Sources: Yeung, 2001; Garrison, 2003; Garrison and Richards, 2004; Garrison, 2005; NMFS, 2004b; Walsh and Garrison, 2006
Table 4.2	Extrapolated (1994-2002) and Observed (2003-2005) Takes and Five Year ITS for Sea Turtles and Sawfish in the HMS Bottom Longline Fishery*. Source: NMFS, 2005, Smith <i>et al.</i> , 2006
Table 4.3	Extrapolated (1999-2002) and Observed (2003-2005) Takes and Five Year ITS for Protected Resources in the Shark Gillnet Fishery*. Source: Carlson, 2003; NMFS, 2003a; Carlson <i>et al.</i> , 2004; Carlson and Bethea, 20054-4
Table 4.4	Percent change in discards of white marlin, blue marlin, sailfish, spearfish, leatherback, loggerhead, and other sea turtles, and bluefin tuna kept and discards combined, based on various time/area closure alternatives without redistribution of effort. + = increase and - = decrease in discards or bycatch. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: Pelagic Observer Program data (2001 – 2003)4-82
Table 4.5	Percent change in discards of white marlin, blue marlin, sailfish, spearfish, leatherback, loggerhead, and bluefin tuna based on various time/area closure alternatives with and without redistribution of effort. + = increase and - = decrease in discards or bycatch. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook data (2001 – 2003)
Table 4.6	Total number of swordfish, bluefin tuna, yellowfin tuna, bigeye tuna, total BAYS (bigeye, albacore, yellowfin and skipjack tuna), reported landed or discarded in the U.S. Atlantic PLL fishery, 1997 – 2003. Pred ¹ = without redistribution of effort, Pred ² = with redistribution of effort. Predictions were predicted back in 2000. Source: HMS Logbook data
Table 4.7	Total number of pelagic sharks, large coastal sharks, dolphin (mahi mahi), and wahoo reported landed or discarded and number of billfish (blue and white marlin, sailfish, spearfish) and sea turtles caught and discarded in the U.S. Atlantic PLL fishery, 1997 – 2003. Pred ¹ = without redistribution of effort, Pred ² = with redistribution of effort. Predictions were predicted back in 2000. Source: HMS logbook data
Table 4.8	Reported distribution of hooks set by area, 1995-2003 (CAR=Caribbean, GOM=Gulf of Mexico, FEC=Florida East Coast, SAB=South Atlantic Bight, MAB=Mid-Atlantic Bight, NEC=Northeast Coastal, NED=Northeast Distant, SAR=Sargasso, NCA=North Central Atlantic, and TUNS=Tuna North & Tuna South). Source: HMS logbook data
Table 4.9	Number of bluefin tuna (BFT), swordfish (SWO), sharks (PEL-pelagic; LCS-Large Coastal Sharks), billfish, and turtles kept and/or discarded in the Mid-Atlantic Bight (MAB) and Northeast Coastal (NEC) areas combined versus all other areas as reported in the pelagic logbook data, 1995-2003. Source: HMS logbook Data
Table 4.10	Alternative B2(a). Example of temporal variation in effectiveness of the closure on discards from May through November with redistribution of effort in (a) all open areas and (b) in the Gulf of Mexico only. Totals and percent changes are for months of May-Nov only. * excluding the NED. += increase and -= decrease in

	discards or bycatch. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook data (2001-2003)
Table 4.11	Alternative B2(a). Percent change in incidental bluefin tuna and target catch and discards with and without redistribution of effort year-round and May-November with redistribution of effort into (a) all open areas and (b) in the Gulf of Mexico only. * excluding the NED + = increase and - = decrease in discards or retained catch. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook data (2001-2003). 4-91
Table 4.12	Alternative B2(b). Percent change in incidental bluefin tuna and target catch and discards without redistribution of effort. * excluding the NED. += increase and -= decrease in discards or retained catch. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook data (2001-2003)
Table 4.13	Alternative B2(c). Percent change in incidental bluefin tuna and target catch and discards without redistribution of effort. * excluding the NED. += increase and -= decrease in discards or retained catch. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook data (2001-2003)
Table 4.14	Alternative B2(b). Example of temporal variation in effectiveness of the closure on discards in June. Totals and percent decreases are for month of June only. * excluding the NED. += increase and -= decrease in discards or bycatch. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook data (2001 – 2003)
Table 4.15	Percent change in kept and discarded catch based on different time/area closure alternatives with and without redistribution of effort = decrease and + = increase in kept and discarded catch; * with redistribution of effort assumes no reduction in the number of hooks set. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook data (2001-2003)
Table 4.16	Alternative B2(c). Example of temporal variation in effectiveness of the time/area closure on discards from April through June. Totals and percent changes are for month of June only. * excluding the NED. $+=$ increase and $-=$ decrease in discards or bycatch. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook data $(2001-2003)$
Table 4.17	Alternative B2(d). Example of temporal variation in effectiveness of the time/area closure on discards. * excluding the NED. $+=$ increase and $-=$ decrease in discards or bycatch. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook data $(2001-2003)$
Table 4.18	Alternative B2(d). Percent change in incidental bluefin tuna and target catch and discards without redistribution of effort. * excluding the NED. += increase and - decrease in discards or retained catch. Three year totals are shown; one year

	averages can be obtained by dividing the three year total by three. Source: HMS Logbook data (2001-2003)
Table 4.19	Alternative B2(e). Example of temporal variation in effectiveness of the time/area
	closure on discards. * excluding the NED. += increase and -= decrease in
	discards or bycatch. Three year totals are shown; one year averages can be
	obtained by dividing the three year total by three. Source: HMS Logbook data
	(2001-2003)4-100
Table 4.20	Alternative B2(e). Percent change in incidental bluefin tuna and target catch and
	discards without redistribution of effort. * excluding the NED. += increase and -
	= decrease in discards or retained catch. Three year totals are shown; one year
	averages can be obtained by dividing the three year total by three. Source: HMS
T 11 401	Logbook data (2001-2003)
Table 4.21	B2(a) and B2(b) Combined. Example of temporal variation in effectiveness of the
	time/area closure on discards for combined alternatives B2(a) and B2(b) year-round, and from May through November for B2(a) and in June for B2(b). *
	excluding the NED. += increase and -= decrease in discards or bycatch. Three
	year totals are shown; one year averages can be obtained by dividing the three
	year total by three. Source: HMS Logbook data (2001-2003)4-102
Table 4.22	B2(a) and B2(b) Combined. Percent change in incidental bluefin tuna and target
	catch and discards without redistribution of effort. * excluding the NED. +=
	increase and - = decrease in discards or retained catch. Three year totals are
	shown; one year averages can be obtained by dividing the three year total by
	three. Source: HMS Logbook data (2001-2003)4-103
Table 4.23	B2(d) and B2(e) Combined. Example of temporal variation in effectiveness of the
	time/area closure on discards for combined alternatives B2(d) and B2(e)year-
	round. * excluding the NED. + = increase and - = decrease in discards or
	bycatch. Three year totals are shown; one year averages can be obtained by
	dividing the three year total by three. Source: HMS Logbook data (2001-2003). 4-
Table 4.24	104 P2(d) and P2(e) Combined Persont aboves in incidental bluefin tune and toward
Table 4.24	B2(d) and B2(e) Combined. Percent change in incidental bluefin tuna and target catch and discards without redistribution of effort. * excluding the NED. +=
	increase and - = decrease in discards or retained catch. Three year totals are
	shown; one year averages can be obtained by dividing the three year total by
	three. Source: HMS Logbook data (2001-2003)4-105
Table 4.25	Alternative B3(a) Charleston Bump modification. Discards of white marlin, blue
14010 1.20	marlin, sailfish, spearfish, leatherback, loggerhead and other sea turtles from the
	portion of the Charleston Bump to remain closed and the portion of the area
	considered for reopening. * excluding the NED. += increase and -= decrease in
	discards or bycatch. Three year totals are shown; one year averages can be
	obtained by dividing the three year total by three. Source: HMS Logbook 1997-
	19994-109
Table 4.26	Alternative B3(b) Northeastern U.S. closure modification. Discards of white
	marlin, blue marlin, sailfish, spearfish, leatherback, loggerhead and other sea
	turtles from the portion of the Northeastern U.S. closure to remain closed and the
	portion of the area considered for reopening. * excluding the NED. + = increase

	and - = decrease in discards or bycatch. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook 1997 - 1999
Table 4.27	Alternative B3(a) Charleston Bump modification. Catches and discards of incidental bluefin tuna and target species in the portion of the area considered for reopening. * excluding the NED. += increase and -= decrease in discards or retained catch. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook 1997-19994-111
Table 4.28	Alternative B3(b) Northeastern U.S. closure modification. Catches and discards of incidental bluefin tuna and target species in the portion of the area considered for reopening. * excluding the NED. += increase and -= decrease in discards or retained catch. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook 1997-19994-112
Table 4.29	Comparison of discards of white marlin, blue marlin, sailfish, spearfish, leatherback and loggerhead sea turtles in the portion of the areas considered for reopening. * excluding the NED. += increase and -= decrease in discards or bycatch. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook 1997-20004-113
Table 4.30	Comparison of of swordfish, incidental bluefin tuna, yellowfin tuna, bigeye tuna, and BAYS in the portion of the areas considered for reopening. * excluding the NED. += increase and -= decrease in discards or retained catch. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook 1997-1999
Table 4.31	Average swordfish lengths (cm LJFL) in the portion of the areas to remain closed and the portion of the areas considered for reopening. The minimum size limit for swordfish is 119 cm LJFL. The mature size is > 180 cm LJFL. Source: Pelagic Observer Program 1992-1999.
Table 4.32	Landings by region from the PLL fishery for swordfish, bluefin tuna, yellowfin tuna and bigeye tuna. The 2004 gross fishing revenues was calculated by converting the 2003 total gross revenues into 2004 dollars (2003 value was multiplied by 1.0266). Source: NMFS, 2004
Table 4.33	Average annual change in gross revenues by species for each of the alternatives in comparison to landings from the Atlantic and Gulf of Mexico. Note: 2003 gross revenues are based on 2003 ex-vessel prices. 2004 gross revenues are based on 2004 ex-vessel prices. The total 2004 annual loss or gain to the fishery was calculated by converting the 2003 total gross revenues into 2004 dollars (2003 value was multiplied by 1.0266) = decrease and + = increase. Source: NMFS, 2005
Table 4.34	Average annual change in gross revenues by species for each of the modifications to the existing time/area closures in comparison to landings from the Atlantic and Gulf of Mexico. Note: 2003 gross revenues are based on 2003 ex-vessel prices. 2004 gross revenues are based on 2004 ex-vessel prices. The total 2004 annual loss or gain to the fishery was calculated by converting the 2003 total gross revenues into 2004 dollars (2003 value was multiplied by 1.0266)= decrease and += increase; * excluding the NED. Source: NMFS, 2005

Table 4.35	Average annual change in gross revenues by species for the B2(a) time/area closures in comparison to landings from the Gulf of Mexico only. Note: 2003 gross revenues are based on 2003 ex-vessel prices. 2004 gross revenues are based on 2004 ex-vessel prices. The total 2004 annual loss or gain to the fishery was calculated by converting the 2003 total gross revenues into 2004 dollars (2003 value was multiplied by 1.0266)= decrease and += increase. Source: NMFS, 2005
Table 4.36	Total number of individual vessels and sets by year in some of the time/area closures. Alternative B4 was not included due to confidentiality concerns. Note: 2004 data is only from January through June 2004. Source: HMS Logbook 2001-2003
Table 4.37	Total number of vessels by state for some of the time/area closures from 2001 through 2003, and 2001 through June of 2004. Alternative B4 was not included due to confidentiality concerns. Source: HMS Logbook 2001-2004 (first six months of 2004)
Table 4.38	Percent of total landings and discards in the Gulf of Mexico (GOM) in comparison to all areas. * excluding the NED. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook 2001-2003
Table 4.39	Percent of total discards in the Gulf of Mexico (GOM) in comparison to all areas (excluding the NED). * excluding the NED. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook 2001-2003
Table 4.40	Finetooth shark landings in lb dw (mt dw) by gear type in the Gulf of Mexico (GOM) and South Atlantic (SA), 1999-2003, as reported in the General Canvass (CN) and Coastal Fisheries (CL) Logbooks. Source: Enric Cortes, pers. comm4-150
Table 4.41	Summary of Small Coastal Shark quotas and landings in commercial and recreational fisheries between 1999-2005. Sources: Data from 1998-2000, Cortés, pers. comm., data from 2001-2004, Cortés, 2005; Carlson and Baremore, 2001; Carlson and Baremore, 2002; Carlson and Baremore, 2003; Carlson <i>et al.</i> , 2004; NMFS, 2005; Cortés and Neer, 2005; Carlson and Bethea, 2006
Table 4.42	Total pounds (lb ww) and percent of total landings by weight of species landed on gillnet trips that landed finetooth sharks, 1999-2004. Species and landings in bold are those currently managed by the HMS Management Division. Species with landings under 10 lbs ww were omitted. Source: Coastal Fisheries Logbook (CFL)
Table 4.43	Total pounds (lb ww) and percent of total landings by weight of species landed on bottom longline trips that landed finetooth sharks, 1999-2004. Species and landings in bold are those currently managed by the HMS Management Division. Species with landings under 10 lbs ww were omitted. Source: CFL4-152
Table 4.44	Number of trips with finetooth shark landings (all gears) by NMFS Statistical Reporting reas, 1999 - 2004. A copy of the map indicating the NMFS Statistical Reporting Areas can be found in Figure 4.18. Bolded areas had more than 20 trips with landings of finetooth sharks. Source: Coastal Fisheries Logbook4-153

Table 4.45	State information on commercial fisheries that may be landing finetooth sharks. 4-155
Table 4.46	Finetooth landings reported from the Florida Trip Ticket program 1999-2004 by area and gear type. All landings in the Exclusive Economic Zone are denoted as EEZ. Source: Steve Brown, Florida Fish and Wildlife Commission Trip Ticket Program
Table 4.47	U.S. Pelagic Longline Dead Discards and Rod & Reel Landings of Atlantic Billfish (MT). <i>Source: Pelagic U.S. National Reports to ICCAT 2003; 2004; 2005</i> 4-160
Table 4.48	U.S. Landings of Atlantic Blue and White Marlin as Reported to ICCAT in Numbers of Fish. Source: U.S. National Reports to ICCAT 2003; 2004; 2005; and Erika Carlsen, NMFS Office of International Affairs, pers. comm4-162
Table 4.49	Tournament Landings and Releases 1999 - 2004 in Number of Fish. Source: NMFS Recreational Billfish Survey (RBS) Database4-162
Table 4.50	MRFSS Estimated Aggregate Landings and Releases of Atlantic Billfish 1999-2004 in Number of Fish (includes Puerto Rico, excludes Texas). Source: NMFS MRFSS Database
Table 4.51	Large Pelagics Survey Estimated Aggregate Landings and Releases of Atlantic Billfish (Virginia to Maine) 1999-2004 in Number of Fish. Source: NMFS LPS Database
Table 4.52	Estimated Post-Release Mortality of White Marlin in Numbers of Fish Based on J-hooks and 35 Percent Post-Release Mortality Rate as derived from Data from the RBS, MRFSS, and LPS. Source: Recreational Billfish Survey; Marine Recreational Fisheries Statistics Survey; and Large Pelagic Survey4-165
Table 4.53	Estimated Post-Release Mortality of Blue Marlin in Numbers of Fish Based on Jhooks and 11 Percent Post-Release Mortality Rate as derived from Data from the RBS, the MRFSS, and LPS. Source: Recreational Billfish Survey; Marine Recreational Fisheries Statistics Survey; and Large Pelagic Survey4-166
Table 4.54	White Marlin Estimated Net Circle Hook Mortality Benefit in Numbers of Fish. Source: RBS, MRFSS, LPS4-170
Table 4.55	Cumulative number of white and blue marlin landed below each minimum size in tournaments from 1999-2004 between the current minimum size for blue (99") and white (66") marlin and the potential increased minimum sizes indicated. A minimum size would be selected between 68-71 inches for WHM and 103-106 for BUM. Source: NMFS RBS Database
Table 4.56	Catch per unit effort and numbers of blue and white marlin kept and released 1999-2004. Source: Billfish Tournament Database, Southeast Fisheries Science Center
Table 4.57	Average Number of Boats per Tournament by Month and State, 1999-2004. Source: Recreational Billfish Survey4-200
Table 4.58	Average Number of Billfish Tournaments by Month and State 1999-2004*. Source: Recreational Billfish Survey
Table 4.59	Alternative F1: Gross revenues associated with the No Action alternative regarding the General Category time-period subquota allocation.*

Table 4.60	Alternative F3(a): Gross revenues associated with the even sub-allocation of the General Category quota between the 8-month fishing season4-217
Table 4.61	Alternative F3(b): Gross revenues associated with the proposed sub-allocation of the General Category quota, providing a BFT winter fishery in the South Atlantic
Table 4.62	Alternative F3(c) (preferred): Gross revenues associated with the proposed sub- allocation of the General Category quota, providing a BFT winter fishery in the South Atlantic
Table 4.63	Alternative F3(d): Gross revenues associated with the proposed sub-allocation of the General Category quota, providing a BFT winter fishery in the South Atlantic. 4-220
Table 4.64	Alternative F8: Gross revenues associated with the proposed rollover cap of 100 percent of the baseline quota allocation
Table 4.65	Fishing years by HMS proposed for Alternatives G1-G34-230
Table 4.66	Average number of commercially harvested BFT by month for 1999-2004. Data source: NERO BFT landings database
Table 4.67	Graphic representation of the distribution of U.S. domestic BFT quota by year. $.4-236$
Table 4.68	Reported Atlantic Commercial Green-stick Gear Catch for 1999-2003*, in Numbers of Fish. Source: PLL Logbook Data
Table 4.69	Reported U.S. Pacific Commercial Green-stick Gear Catch for 2002-2004, in Numbers of Fish and Weight in Pounds. Source: State of Hawaii, 20064-243
Table 4.70	Handline catches, in Numbers of Fish, for 2000 – 2004. Source: HMS Logbook. Note that confidential data cannot be released and are marked by an *4-246
Table 4.71	List of "Indicator" Species to Determine Composition of Catch4-261
Table 4.72	Historical Per Trip Average Catch Composition of "Indicator" Species. Source: NMFS Coastal Logbook
Table 4.73	Historical Per Trip Average Catch Composition of "Indicator" Species. Source: Pelagic Longline Logbook
Table 4.74	Numbers of Trips Reported as Using Handline Gear in the HMS Logbook and Numbers of Those Trips that were "Handline-Only." Source: HMS Logbook. Note that confidential data cannot be released and are marked by an *4-274
Table 4.75	Impacts of alternatives considered. The symbols +, -, and 0 refer to positive, negative, and zero impacts respectively. The expected impacts should be compared to other alternatives within that issue, not to the impacts between issues. See preceding section for details of impacts of each alternative4-308
Table 6.1	Estimates of the total ex-vessel annual revenues of Atlantic HMS fisheries. Sources: NMFS 2005a; Cortes, 2005; and bluefin tuna dealer reports from the Northeast Regional Office. 6-2
Table 7.1	Summary of the Net Economic Benefits and Costs of Alternatives7-4
Table 9.1	Demographic Profile of the United States. Source: U.S. Census, 1990 and 2000.
Table 9.2	Maine Demographic Profile. Source: U.S. Census, 1990 and 20009-8
Table 9.3	New Hampshire Demographic Profile. Source: U.S. Census, 1990 and 2000 9-9

Table 9.4	Massachusetts Demographic Profile. Source: U.S. Census, 1990 and 2000 9-10
Table 9.5	Demographic Profile of Gloucester, Massachusetts. Source: U.S. Census, 1990 and 2000
Table 9.6	Demographic Profile of New Bedford, Massachusetts. Source: U.S. Census, 1990 and 2000
Table 9.7	Rhode Island Demographic Profile. Source: U.S. Census, 1990 and 2000 9-18
Table 9.8	Connecticut Demographic Profile. Source: U.S. Census, 1990 and 20009-19
Table 9.9	New York Demographic Profile. Source: U.S. Census, 1990 and 20009-20
Table 9.10	New Jersey Demographic Profile. Source: U.S. Census, 1990 and 20009-22
Table 9.11	Demographic Profile of Barnegat Light. Source: U.S. Census, 1990 and 20009-23
Table 9.12	Demographic Profile of Brielle, New Jersey. Source: U.S. Census, 1990 and 20009-26
Table 9.13	Delaware Demographic Profile. Source: U.S. Census, 1990 and 20009-28
Table 9.14	Maryland Demographic Profile. Source: U.S. Census, 1990 and 20009-29
Table 9.15	Virginia Demographic Profile. Source: U.S. Census, 1990 and 20009-31
Table 9.16	Demographic Profile of North Carolina. Source: U.S. Census, 1990 and 2000 9-32
Table 9.17	Demographic Profile of Hatteras, North Carolina Source: U.S. Census, 1990 and 20009-34
Table 9.18	Demographic Profile of Wanchese, North Carolina. Source: U.S. Census 1990 and 20009-37
Table 9.19	South Carolina Demographic Profile. Source: U.S. Census, 1990 and 2000 9-39
Table 9.20	Georgia Demographic Profile. Source: U.S. Census, 1990 and 20009-40
Table 9.21	Florida Demographic Profile. Source: U.S. Census, 1990 and 20009-42
Table 9.22	Demographic Profile of Pompano Beach, Florida. Source: U.S. Census 1990 and 2000
Table 9.23	Demographics of Fort Pierce, Florida. Source: U.S. Census 1990 and 20009-45
Table 9.24	Demographic Profile for Madeira Beach, Florida. Source: U.S. Census 1990 and 20009-47
Table 9.25	Demographic Profile for Panama City, Florida. Source: U.S. Census 1990 and 20009-50
Table 9.26	Demographic Profile for Islamorada, Florida. Source: U.S. Census, 1990 and 2000
Table 9.27	Alabama Demographic Profile. Source: U.S. Census, 1990 and 20009-54
Table 9.28	Mississippi Demographic Profile. Source: U.S. Census, 1990 and 20009-55
Table 9.29	Louisiana Demographic Profile. Source: U.S. Census, 1990 and 2000 9-56
Table 9.30	Demographic Profile of Venice, Louisiana. Source: U.S. Census, 1990 and 2000.
Table 9.31	Demographic Profile of Dulac, Louisiana. Source: U.S. Census, 1990 and 2000.9-63
Table 9.32	Texas Demographic Profile. Source: U.S. Census, 1990 and 20009-65

Table 9.33	Puerto Rico Demographic Profile. Source: U.S. Bureau of the Census, 1990 and 2000
Table 9.34	Number and Percentage of HMS Angling Permits by State and Country in 2005.
Table 9.35	Number and Percentage of HMS Charter/Headboat Permits by State and Country in 2005
Table 9.36	Number and Percentage of Commercial Tuna Permits by State and Country in 20059-72
Table 9.37	Number and Percentage of HMS Dealers by State and Country as of February 2006 (sharks and swordfish) and for calendar year 2005 (tunas)73
Table 9.38	Number and Percentage of Directed and Incidental Shark Permit Holders by State as of February 2006
Table 9.39	Number and Percentage of Swordfish Permit Holders by State as of February 20069-75
Table 10.1.	Impact assessment of HMS fishing gear on HMS and non-HMS EFH. '-' indicates that the gear type is not used in these habitat types. Habitat impacts are as follows: negligible = 0, low = +, medium = ++, high = +++, unknown=?, and a blank indicates not evaluated. Source: Symbols before the slash are from the Caribbean FEIS, 2004 (Table 3.15a). The symbols after the slash are taken from Barnette, 2001
Table 10.2.	A comprehensive list of all gear types used in HMS fisheries10-28
Table 10.3.	Fishing gear types regulated in Federal waters of the Northeast region and their effects on HMS EFH. Habitat impact is as follows: minimal/negligible = 0. Source: Stevenson <i>et al.</i> , 2004
Table 10.4.	Fishing gear types regulated in Federal waters in the Southeast region and their effects on HMS EFH. Habitat impact is as follows: minimal/negligible = 0. Source: Hamilton, 2000; Barnette, 2001; GOMFMC FEIS 2004
Table 10.5.	Non-FMP Fishing Gear in the Northeast region and their effects on HMS EFH. Habitat impact is as follows: minimal/negligible = 0. Source: Stevenson <i>et al.</i> , 2004
Table 10.6.	Non-FMP Fishing Gear in Southeast region: Effects of other fishing gear on HMS EFH. Habitat impact is as follows: minimal/negligible = 0. Source: Hamilton, 2000; Barnette, 2001; GOMFMC, 2004
Table A.1	The decrease (-) or increase (+) in the number of discards of white marlin, blue marlin, sailfish, spearfish, leatherback and loggerhead sea turtles and bluefin tuna based on the various time/area closures. * excluding NED. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS logbook data 2001-2003
Table A.2	Percent reduction (-) or increase (+) in discards of white marlin, blue marlin, sailfish, spearfish, leatherback and loggerhead sea turtles and bluefin tuna based on the various time/area closure alternatives with and without redistribution of effort. (* = was not analyzed). Source: HMS Logbook data (2001-2003) A-10
Table A.3	The decrease (-) or increase (+) in the number of each retained species caught or discarded based on the various time/area closure alternatives without

	redistribution of effort. *excluding the NED. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook data (2001-2003)
Table A.4	Percent reduction (-) or increase (+) in the retained catch and discards based on the various time/area closure alternatives without redistribution of effort. Source: HMS Logbook data (2001-2003)
Table A.5	The decrease (-) or increase (+) in the number of each retained species caught or discarded based on the various time/area closure alternatives with redistribution of effort. (* = was not calculated). ¹ excluding the NED. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook data (2001-2003)
Table A.6	Percent reduction (-) or increase (+) in the retained catch and discards based on the various time/area closure alternatives with redistribution of effort. (* = was not calculated). Source: HMS Logbook data (2001-2003)
Table A.7	Percent reduction (-) or increase (+) in the number of hooks set; discards of white marlin, blue marlin, sailfish, spearfish, leatherback, loggerhead, and other sea turtles based on various time/area closure alternatives without redistribution of effort. Source: HMS Logbook data (2001-2003)
Table A.8	Percent reduction (-) or increase (+) in discards of white marlin, blue marlin, sailfish, spearfish, leatherback, loggerhead, and other sea turtles, and bluefin tuna kept and discards combined, based on various time/area closure alternatives without redistribution of effort. Source: Pelagic Observer Program data (2001-2003).
Table A.9	Alternative B2(f). Temporal variation in effectiveness of Gulf of Mexico time/area closure in terms of percent reduction (-) or increase (+) of white marlin, blue marlin, sailfish, spearfish, leatherback, loggerhead, and bluefin tuna discards. A negative sign indicates an increase in bycatch. *excluding the NED. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook data (2001-2003)
Table A.10	Alternative B2(g). Temporal variation in effectiveness of the Northeast time/area closure from June through October in terms of percent reduction (-) or increase (+) of white marlin, blue marlin, sailfish, spearfish, leatherback, loggerhead, and other sea turtle discards. *excluding the NED. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook data (2001-2003)
Table A.11	Alternative B2(h). Temporal variation in effectiveness of the Southeast time/area closure closure in terms of percent reduction (-) or increase (+) of white marlin, blue marlin, sailfish, spearfish, leatherback, loggerhead, and other sea turtle discards. *excluding the NED. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook data (2001-2003)
Table A.12	Alternative B2(i). Temporal variation in effectiveness of the closure on the east coast of Florida in terms of percent reduction (-) or increase (+) of white marlin, blue marlin, sailfish, spearfish, leatherback, loggerhead, and other sea turtle discards. *excluding the NED. Three year totals are shown; one year averages

	can be obtained by dividing the three year total by three. Source: HMS Logbook data (2001-2003)
Table A.13	Alternative B2(j). Temporal variation in effectiveness of the Gulf of Mexico time/area closure in terms of percent reduction (-) or increase (+) of white marlin, blue marlin, sailfish, spearfish, leatherback, loggerhead, and other sea turtle discards. *excluding the NED. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook data (2001-2003)
Table A.14	Alternative B2(k). Temporal variation in effectiveness of the Caribbean time/area closure in terms of percent reduction (-) or increase (+) of white marlin, blue marlin, sailfish, spearfish, leatherback, loggerhead, and other sea turtle discards. Landings were only reported for the four months listed. *excluding the NED. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook data (2001-2003)
Table A.15	Alternative B2(f). Temporal variation in effectiveness of the Gulf of Mexico time/area closure in terms of percent reduction (-) in discards and retained catch. *excluding the NED. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook data (2001-2003)
Table A.16	Alternative B2(g). Temporal variation in effectiveness of the Northeast time/area closure from June through October closure in terms of percent reduction (-) in discards and retained catch. *excluding the NED. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook data (2001-2003)
Table A.17	Alternative B2(h). Temporal variation in effectiveness of the time/area closure from March through November closure in terms of percent reduction (-) in discards and retained catch. *excluding the NED. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook data (2001-2003)
Table A.18	Alternative B2(i). Temporal variation in effectiveness of the closure of the east Florida in terms of percent reduction (-) in discards and retained catch. *excluding the NED. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook data (2001-2003)
Table A.19	Alternative B2(j). Temporal variation in effectiveness of the time/area closure in the Gulf of Mexico in terms of percent reduction (-) in discards and retained catch. *excluding the NED. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook data (2001-2003)
Table A.20	Alternative B2(k). Temporal variation in effectiveness of the time/area closure in the Caribbean in terms of percent reduction (-) in discards and retained catch. *excluding the NED. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook data (2001-2003)

Table A.21	Swordfish lengths (cm lower jaw fork length; LJFL) in the portion of the areas to remain closed and the portion of the areas considered for reopening. The minimum size limit for swordfish is 119 cm LJFL. The mature size is > 180 cm LJFL. Significant differences are shaded. Source: Pelagic Observer Program 1992-1999
Table A.22	Alternative B3(c) Florida East Coast modification. Discards of white marlin, blue marlin, sailfish, spearfish, leatherback, loggerhead and other sea turtles in the portion of the area to remain closed and the portion of the area considered for reopening. *excluding the NED. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook 1997-1999
Table A.23	Alternative B3(d) Desoto Canyon modification. Discards of white marlin, blue marlin, sailfish, spearfish, leatherback, loggerhead and other sea turtles in the portion of the area to remain closed and in the portion of the area considered for reopening. Source HMS Logbook 1997-1999. *excluding the NED. Three year totals are shown; one year averages can be obtained by dividing the three year total by three
Table A.24	Comparison of discards of white marlin, blue marlin, sailfish, spearfish, leatherback and loggerhead sea turtles in the portion of the areas considered for reopening= decrease += increase. *excluding the NED. Four year totals are shown; one year averages can be obtained by dividing the four year total by four. Source: HMS Logbook 1997-2000
Table A.25	Comparison of catch of swordfish, bluefin tuna, yellowfin tuna, bigeye tuna, and BAYS in the portion of the areas considered for reopening. *excluding the NED. Four year totals are shown; one year averages can be obtained by dividing the four year total by four. Source: HMS Logbook 1997-2000
Table A.26	Alternative B3(c) modification of East Florida Coast time/area closure. Catch and discards of various species in the portion of the area to remain closed and in the portion of the area considered for reopening. *excluding the NED. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook data (1997-1999)
Table A.27	Alternative B3(d) modification of the DeSoto Canyon time/area closure. Catch and discards of various species in the portion of the area to remain closed and in the portion of the area considered for reopening. *excluding the NED. Three year totals are shown; one year averages can be obtained by dividing the three year total by three. Source: HMS Logbook data (1997-1999)
Table A.28.	An example of the calculations for the redistribution of fishing effort model. This example calculates the number of discards of loggerhead sea turtles considering the redistribution of fishing effort for alternative B2(d). A negative sign indicates an increase in discards
Table A.29	An example of how the modified redistribution of fishing effort was calculated. This example calculates the number of discards of white marlin considering the redistribution of fishing effort in the Gulf of Mexico and area 6 only for alternative B2(c) from April through June. A negative sign indicates an increase in discards, and a positive value indicates a decrease in discards

Table A.30	An example of how the modified redistribution of fishing effort was calculated. This example calculates the number of discards of white marlin considering the redistribution of fishing effort in the Gulf of Mexico and area 6 only for alternative B2(c) from April through June. A negative sign indicates an increase in discards, and a positive value indicates a decrease in discards
Table A.31	Comparison between the percent change of bycatch for different closures for 2001-2003 data (used in the draft Consolidated FMP) and 2001-2004 (first six months of 2004 included) without redistribution of effort. A negative (-) sign indicates a reduction in bycatch. Source: HMS Logbook data 2001-2004 (first six months of 2004)
Table A.32	Comparison in the number of hooks, discards, and CPUEs (# discards/# hooks in a particular area) between July through December of 2001-2003 and 2004 data. The yearly averages for the 6 months in 2001-2003 for CPUEs, hooks, and discards are shown in parentheses. Data source: HMS Logbook data July through December 2001-2003 and 2004
Table A.33	Comparison of the number of hooks, discards, species kept, and CPUEs (#discards/#hooks or #kept/#hooks in a particular area) between July through Dec. of 2001-2003 and 2004 data. The yearly averages for the 6 months in 2001-2003 for CPUEs, hooks, discards, and species kept are shown in parentheses Data source: HMS Logbook data July through December 2001-2003 and 2004. A-46
Table A.34	Percent change in reported landings by area from July through December where: a) 2001-03 vs. 1997-99; b) 2004 vs. 2001-03; and c) 2004 vs. 1997-99 (1997-99 and 2001-03 are mean reported landings). Source: HMS Logbook data A-48
Table A.35	Percent change in reported landings by area from July through December where: a) 2001-03 vs. 1997-99; b) 2004 vs. 2001-03; and c) 2004 vs. 1997-99 (1997-99 and 2001-03 are mean reported landings). Source: HMS Logbook data A-50
Table A.36	Characteristics of vessels fishing in the Gulf of Mexico and vessels moving out of Gulf of Mexico
Table A.37	Alternative B2(a) May through November. Cumulative number kept and discarded (over 3 1/2 years) with and without redistribution of effort in the Gulf of Mexico and Area 6 combined. Minus sign indicates a decrease. *excluding the NED. Three and one half year totals are shown; one year averages can be obtained by dividing the totals by 3.5. Data source: HMS Logbook data 2001-2004 (first six months of 2004)
Table A.38	Alternative B2(b) June only. Cumulative number of discards (over 3 1/2 years) with and without redistribution of effort in the Atlantic only. Minus signs indicate a decrease. *excluding the NED. Three and one half year totals are shown; one year averages can be obtained by dividing the totals by 3.5. Data source: HMS Logbook data 2001-2004 (first six months of 2004)
Table A.39	Alternative B2(b) June only. Cumulative number kept and discarded (over 3 1/2 years) with and without redistribution of effort in the Atlantic only. Minus signs indicate a decrease. *excluding the NED. Three and one half year totals are shown; one year averages can be obtained by dividing the totals by 3.5. Data source: HMS Logbook data 2001-2004 (first six months of 2004)

Table A.40	Alternative B2(c) BFT Petition April through June. Cumulative number of discards (over 3 1/2 years) with and without redistribution of effort in the Gulf of Mexico and Area 6 combined. Minus signs indicate a decrease. *excluding the NED. Three and one half year totals are shown; one year averages can be obtained by dividing the totals by 3.5. Data source: HMS Logbook data 2001 - 2004 (first six months of 2004)
Table A.41	Alternative B2(c) BFT Petition April through June. Cumulative number of kept and discarded (over 3 1/2 years) species with and without redistribution of effort in the Gulf of Mexico and Area 6 combined. Minus sign indicates a decrease. *excluding the NED. Three and one half year totals are shown; one year averages can be obtained by dividing the totals by 3.5. Data source: HMS Logbook data 2001-2004 (first six months of 2004)
Table B.1	1999 FMP size ranges for different life stages of sharks
Table B.2	Size ranges used in this Amendment for mapping distribution data for different life stages of sharks
Table B.3	Blacktip shark (<i>Carcharinus limbatus</i>) Life History and Habitat Characteristics. From Amendment 1 to the FMP
Table B.4	Dusky shark (<i>Carcharinus obscurus</i>) Life History and Habitat Characteristics B-66
Table B.5	Sandbar shark (<i>Carcharinus plumbeus</i>) Life History and Habitat CharacteristicsB-67
Table B.6	Nurse shark (<i>Ginglymostoma cirratum</i>) Life History and Habitat Characteristics
Table B.7	Essential fish habitat maps by species B-70
Table B.8	List of abbreviations and acronyms for EFH data sources used in the maps B-71
Table C.1	Estimated Domestic Aggregate Pelagic Longline Atlantic White Marlin Mortalities in Numbers of Fish 2001-2004. Source: Pelagic Longline Logbook; Kerstetter, 2006
Table C.2	Estimated Domestic Aggregate Recreational Atlantic White Marlin Mortalities, in Number of Fish, as Derived from the RBS Database by Combining Retained Fish and Dead Discarded Fish with Estimated Post-Release Mortalities (PRM) (applying a 35% post-release mortality estimate) 2001-2004. Source: Recreational Billfish Survey; Horodysky, 2005
Table C.3	Estimated Domestic Aggregate Recreational Atlantic White Marlin Mortalities, in Number of Fish, as Derived from the MRFSS Database by Combining Retained Fish and Dead Discarded Fish with Estimated Post-Release Mortalities (PRM) (applying a 35% post-release mortality estimate) 2001-2004. Source: Marine Recreational Fishing Statistics Survey; Horodysky, 2005
Table C.4	Estimated Domestic Aggregate Recreational Atlantic White Marlin Mortalities, in Number of Fish, as Derived from the LPS Database by Combining Retained Fish and Dead Discarded Fish with Estimated Post-Release Mortalities (PRM) (applying a 35% post-release mortality estimate) 2001-2004. Source: Large Pelagics Survey; Horodysky, 2005; Large Pelagic Survey; Horodysky, 2005 C-4

MASTER LIST OF FIGURES

Figure 2.1	Existing time/area closures in HMS fisheries. Inset shows extent of the Northeast Distant restricted fishing area. All closures except the Mid-Atlantic are applicable to pelagic longline gear only. The Mid-Atlantic Closure is applicable to bottom longline gear only. Note: the Northeast Distant (NED) was a closed area to all vessels as of 2001. It became the NED Restricted Fishing Area on June 30, 2004 when it was opened to those participating in the NED experiment2-19
Figure 2.2	Map showing areas being considered for new time/area closures to reduce non-target HMS and protected species interactions2-20
Figure 2.3	Map showing areas considered for modifications to existing closures. Note: only alternatives B3(a) and (b) were further analyzed2-21
Figure 2.4	Pelagic and Bottom Longline Sets in the Madison-Swanson (upper left) and Steamboat Lumps (lower right) Marine Reserves. Note: one set for the Commercial Shark Fishery Observer Program (CSFOP) was in 2005. Although not indicated, no new sets were recorded for the CSFOP in 2004. Source: HMS Logbook, Pelagic Observer Program, and CSFOP. The Desoto Canyon closure is also shown for reference
Figure 2.5	Map showing the potential closed area to bottom longline gear to reduce bycatch of endangered smalltooth sawfish. Grey dots are locations of observed bottom longline sets. Source: CSFOP 1994-20062-23
Figure 2.6	Map showing time/area closure alternatives considered but not further analyzed at this time to reduce white marlin and other protected species interactions 2-24
Figure 2.7	The New York Bight set-aside is defined as an area comprising the waters South and West of a straight line originating at a point on the southern shore of Long Island, NY, at 72° 27' W. Long. (Shinnecock Inlet) and running South southeast 150° true, and north of 38° 47' N. Lat
Figure 2.8	Alternative F1: No Action. Suballocation of the BFT General Category Quota among the current three time-periods. New York Bight set-aside is subtracted from the General Category quota and then the time-period allocations are determined
Figure 2.9	The Angling category North/South dividing line, located at 39° 18' N. Lat. (Great Egg Inlet, NJ)
Figure 2.10	Alternative F3a: Equal General category subquota allocation percentages for each month of the BFT fishing season2-38
Figure 2.11	Alternative F3b: Proposed General category time-period subquota allocation percentages
Figure 2.12	Alternative F3c: Proposed General category time-period subquota allocation percentages
Figure 2.13	Alternative F3d: Proposed General category time-period subquota allocation percentages embodying the NCDMF Petition for Rulemaking2-40
Figure 2.14	Alternative F5: No Action. U.S. BFT Domestic Quota Category Allocation Percentages
Figure 2.15	A Diagram of a Typical Speargun Fishing Gear Configuration (courtesy of Matthew Richards)2-51

Figure 2.16	A Diagram of the Recreational Configuration of Green-stick Fishing Gear. Source: Wescott, 1996
Figure 2.17	A Diagram of the Commercial Configuration of Green-stick Fishing Gear. Source: Wescott, 1996
Figure 2.18	A Diagram of a Buoy Gear with Four Floatation Devices Attached (courtesy of Dave Meyer)2-57
Figure 3.1	Illustration of the status determination criteria and rebuilding terms 3-41
Figure 3.2	Reported catches (mt whole weight) of Atlantic Swordfish, including discards for 1950-2004. Source: SCRS, 2005
Figure 3.3	Estimated fishing mortality rate relative to FMSY (F/FMSY) for the period 1959-2001 (median with 80 percent confidence bounds based on bootstrapping are shown). Source: SCRS 2004
Figure 3.4	Annual yield (mt) (whole weight) for North Atlantic swordfish relative to the estimated MSY level. Source: SCRS 2004
Figure 3.5	Western Atlantic bluefin tuna spawning biomass (t), recruitment (numbers) and fishing mortality rates for fish of age 8+, estimated by the Base Case VPA run. Source: ICCAT, 2004
Figure 3.6	Abundance indices in numbers of BET. All ages are aggregated. Source: ICCAT, 2004
Figure 3.7	Trajectory of the BET biomass modeled in production model analysis (middle line) bounded by upper and lower lines denoting 80 percent confidence intervals. Source: ICCAT, 2004
Figure 3.8	Comparison of relative biomass trends calculated using VPA and non-equilibrium production models. Source: ICCAT, 2004
Figure 3.9	Comparison of relative fishing mortality trends calculated using VPA and non-equilibrium production models. Source: ICCAT, 20043-71
Figure 3.10	North Atlantic albacore spawning stock biomass and recruits with 80 percent confidence limits. Source: ICCAT, 2004
Figure 3.11	Estimated catches (including landings and dead discards in mt) of blue marlin in the Atlantic by region. The 2003 catch reported to ICCAT is preliminary and is not included in this figure. Weights are in metric tones, whole weight. Source: SCRS, 2005
Figure 3.12	Composite CPUE series (symbols) used in the blue marlin assessment compared to model estimated median relative biomass (solid lines) from bootstrap results (80 percent confidence bounds shown by dotted lines). Source: SCRS, 20053-88
Figure 3.13	Estimated median relative fishing mortality trajectory for Atlantic blue marlin (center, dark line) with approximate 80 percent confidence range (light lines) obtained from bootstrapping. Source: SCRS, 2005
Figure 3.14	Geographical distribution of reported catches of blue marlin for the period 1956-2003. Source: SCRS, 2005
Figure 3.15	Estimated catches (including landings and dead discards in t) of blue marlin in the Atlantic by region (1950-2004). Source: SCRS, 2005
Figure 3.16	Estimated biomass ratio B_{2000}/B_{MSY} (solid line, no symbols) and fishing mortality ratio F2000/FMSY (solid line with symbols) from the production model fitted to

	the continuity case for white marlin. Ratios of last three years have been adjusted for retrospective pattern. Broken lines show unadjusted ratios. Note that scales are different for each ratio. Source: SCRS, 2004
Figure 3.17	Geographical distribution of white marlin catches for the period 1956-2003. Source: SCRS, 2005
Figure 3.18	Reported catch of white marlin (Task I) in the North and South Atlantic for longline (LL) gear and other (OTH) gears. Source: SCRS, 20053-96
Figure 3.19	Evolution of estimated sailfish/spearfish catches in the Atlantic (landings and dead discards, reported and carried over) in the ICCAT Task I database during 1956-2002 for the east and west stocks. The 2003 catch reported to ICCAT is preliminary and is not included in this figure. Weights are in metric tons, whole weight. Source: SCRS, 2005
Figure 3.20	Available standardized CPUE for western Atlantic sailfish/spearfish for the period 1967-2000, including Japanese, U.S., and Venezuelan time series data. Source: SCRS, 2005
Figure 3.21	Estimated sailfish "only" catches based on the new procedure for splitting combined sailfish and longbill spearfish catches from 1956-2000. Weights are in metric tons, whole weight. Source: SCRS, 2005
Figure 3.22	Geographical distribution of sailfish/spearfish catches between 1950-2003. Source: SCRS, 2005
Figure 3.23	Evolution of estimated sailfish/spearfish catches in the Atlantic (landings and dead discards, reported and carried over) in the ICCAT Task I database during 1956-2004 for the east and west stocks. Source: SCRS, 2005
Figure 3.24	Estimated spearfish "only" catches in the Atlantic based on the new procedure for splitting combined sailfish and spearfish catches from 1956-2000. Weights are in metric tons, whole weight. Source: SCRS, 2005
Figure 3.25	Typical U.S. Pelagic Longline Gear. Source: Arocha, 1996
Figure 3.26	Different Pelagic Longline Gear Deployment Techniques. Source: Hawaii Longline Association and Honolulu Advertiser
Figure 3.27	Geographic Areas Used in Summaries of Pelagic Logbook Data. Source: Cramer and Adams, 2000
Figure 3.28	Areas Closed to Pelagic Longline Fishing by U.S. Flagged Vessels3-162
Figure 3.29	Distribution of Atlantic Longline Catches for all Countries 1990-1999. Source: SCRS, 2004
Figure 3.30	Total Number of Swordfish Caught, Kept and Released in Three Sampled Recreational Swordfish Tournaments off Southeast Florida during 2002. Source: J. Levesque, pers. comm
Figure 3.31	Observed sea turtle interactions and observed sets (smaller grey circles) in the shark bottom longline fishery from 1994-2004. Source: Burgess and Morgan, 2004
Figure 3.32	Observed sawfish interactions and observed sets (smaller grey circles) in the shark bottom longline fishery from 1994-2006. Source: Burgess and Morgan, 2004

Figure 3.33	Average Annual Yen/\$ Exchange Rate and Average U.S. BFT Ex-vessel \$/lb (dw) for all gears: 1971-2003. Source: Federal Reserve Bank (www.stls.frb.org) and Northeast Regional Office
Figure 4.1	Existing time/area closures in HMS fisheries. Inset shows extent of the Northeast Distant restricted fishing area. All closures except the Mid-Atlantic are applicable to PLL gear only. The Mid-Atlantic Closure is applicable to bottom longline gear only. Note: the Northeast Distant (NED) was a closed area to all vessels as of 2001. It became the NED Restricted Fishing Area on June 30, 2004 when it was opened to those participating in the NED experiment
Figure 4.2	Map showing areas being considered for new time/area closures to reduce non-target HMS and protected species interactions4-70
Figure 4.3	Map showing areas being considered for modifications to existing closures4-71
Figure 4.4	Pelagic and Bottom Longline Sets in the Madison-Swanson (upper left) and Steamboat Lumps (lower right) Marine Reserves. Note: one set for the CSFOP was in 2005. Although not indicated, no new sets were recorded for the CSFOP in 2004. Source: HMS Logbook, Pelagic Observer Program, Shark Observer Program. The Desoto Canyon closure is also shown for reference
Figure 4.5	Map showing the potential closed area to bottom longline gear to reduce bycatch of endangered smalltooth sawfish. Grey dots are locations of observed bottom longline sets. Source: CSFOP 1994 – 2006
Figure 4.6	Map showing all reported sets and white marlin interactions: a) shows the number of interactions in absolute numbers, b) shows CPUE (per 1,000 hooks). Source: HMS Logbook 2001-20034-74
Figure 4.7	Map showing all observed sets and white marlin interactions. Source: Pelagic Observer Program 2001-20034-75
Figure 4.8	Map showing all reported sets and bluefin tuna discards: a) shows the number of interactions in absolute numbers, b) shows CPUE (per 1,000 hooks). Source: HMS Logbook 2001-2003
Figure 4.9	Map showing all observed sets and bluefin tuna discards. Source: Pelagic Observer Program 2001-20034-77
Figure 4.10	Map showing all reported sets and leatherback sea turtle interactions: a) shows the number of interactions in absolute numbers, b) shows CPUE (per 1,000 hooks). Source: HMS Logbook 2001-20034-78
Figure 4.11	Map showing all reported sets and loggerhead sea turtle interactions: a) shows the number of interactions in absolute numbers, b) shows CPUE (per 1,000 hooks). Source: HMS Logbook 2001-20034-79
Figure 4.12	Map showing all observed sets and sea turtle interactions. Source: Pelagic Observer Program 2001 – 2003
Figure 4.13	Map showing discards of bluefin tuna in the Atlantic and Gulf of Mexico from 2001-2003. The large box is not a proposed time/area closure, but is shown for illustrative purpose only to delineate an area with high bluefin tuna discards. The bluefin tuna discards are listed by month below the box. Source: HMS Logbook 2001-2003

Figure 4.14	Map showing areas being considered for modifications to existing closures and juvenile swordfish data (<180 cm LJFL). The minimum size limit for swordfish is 119 cm LJFL. Source: Pelagic Observer Program 1997-19994-106
Figure 4.15	Map showing areas being considered for modifications to existing closures and adult swordfish data (>180 cm LJFL) from the Pelagic Observer Program. Source: Pelagic Observer Program 1997-1999
Figure 4.16	Map showing modification to the existing Northeast United States closure and bluefin tuna discards. Source: HMS Logbook 1997-19994-108
Figure 4.17	Map showing smalltooth sawfish bycatch. Numbers in parentheses in the legend are the number of sawfish interactions. Gray dots indicate the location of each observed bottom longline set. Source: CSFOP 1994-20064-116
Figure 4.18	NMFS Statistical Reporting Areas. Areas 3, 23, 2780, 2880, 2980, 3079, and 3379 all had greater than 20 trips that reported finetooth shark landings in the Coastal Fisheries Logbook between 1999-2004 as indicated in Table 4.44 Number of trips with finetooth shark landings (all gears) by NMFS Statistical Reporting reas, 1999 - 2004. A copy of the map indicating the NMFS Statistical Reporting Areas can be found in Figure 4.18. Bolded areas had more than 20 trips with landings of finetooth sharks. Source: Coastal Fisheries Logbook4-154
Figure 4.19	Finetooth shark trips by month and gear type between 1999-2004. E/H = Electric Reel, Bandit Gear, Handline; GN = Gillnet; and LL = Longline. Source: Coastal Fisheries Logbook
Figure 4.20	U.S. Pelagic Longline Dead Discards and Rod & Reel Landings of Atlantic Billfish. Source: U.S. National Reports to ICCAT 2003; 2004; 20054-160
Figure 4.21	Number of blue marlin landed in tournaments between 1999-2004 by size (inches, LJFL). Source: NMFS RBS database
Figure 4.22	Number of white marlin landed in tournaments between 1999-2004 by length (inches, LJFL). Source: NMFS RBS Database
Figure 4.23	Map of the East Florida Coast closed area (solid shaded area) and the boundary of the U.S. EEZ (thin line wrapping around the coast). The inset is a close-up of the closed area depicting both the old and proposed (new) boundary coordinates. The small grey dots represent locations of longline sets from the year 2000 through the first half of 2004
Figure 9.1	Location of HMS Angling Permit Holders in 2005 and the percentage of Angling permit holders for the top five states
Figure 9.2	Location of HMS Angling Permit Holders in 2005 by region9-70
Figure 9.3	Location of the HMS Charter/Headboat Permit Holders in 2005 and the percentage of HMS Charter/Headboat permit holders for the top five states9-71
Figure 9.4	Location of the Commercial Tuna Permit Holders in 2005 (all gear categories - harpoon, longline, purse seine, and trap) and the percentage of commercial tuna permit holders for the top five states
Figure 9.5	Location of all HMS Dealer Permit Holders as of February 2006 for shark and swordfish permits and for fishery year 2005 for tunas and the percentage of total HMS dealer permit holders for the top five states

Figure 9.6	Location of the Shark Directed and Incidental Permit Holders as of February 2006 and percentage of shark permit holders for the top five states			
Figure 9.7	Location of the Swordfish Permit Holders as of February 2006 and the percentage of swordfish permit holders for the top five states			
Figure A.1	Map showing time/area closure alternatives considered but not further analyzed at this time (see Section 2.1.2) to reduce white marlin and other protected species interactions			
Figure A.2	Map showing areas considered for modifications to existing closures. Note: only alternatives B3(a) and (b) were further analyzed			
Figure A.3	Map showing areas considered for modifications to existing closures and juvenile swordfish data (<180 cm LJFL). The minimum size limit for swordfish is 119 cm LJFL. Note: only alternatives B3(a) and (b) were further analyzed. Source Pelagic Observer Program 1997-1999			
Figure A.4	Map showing areas considered for modifications to existing closures and adult swordfish data from the Pelagic Observer Program. Note: only alternatives B2(a) and (b) were further analyzed. Source Pelagic Observer Program 1997-1999. A-32			
Figure A.5	Map showing the different areas that were used in the fleet mobility analysis. A-52			
Figure A.6	Map showing the number of vessels fishing in different areas with their respective homeports listed. "FL east" signifies that a vessel's homeport was in the east coast of Florida. "FL west" signifies that a vessel's homeport was in the west coast of FL, and "FL east/west" signifies that the vessel's homeport was in the Florida Keys			
Figure A.7	Map showing vessels originating from the Gulf of Mexico and fishing in Atlantic Areas 2B through 6. The percentage of hooks represents the percentage of hooks that moved out of the Gulf of Mexico and into Areas 2B, 3, 4, 5, and 6			
Figure A.8	Map of pelagic longline sets within and outside of the U. S. EEZ. a) extent of pelagic longline sets inside and outside U.S. EEZ, b) inset shows the size of B2(a) and B2(c) relative to the scope of pelagic longline sets inside and outside of U. S. EEZ, c) the distance pelagic longline vessels have made relative to the U. S. coastline, d) the distance it would take a vessel to travel from the Gulf of Mexico to the Mid-Atlantic Bight area. Source: HMS logbook data from January 2001 to June 2004.			
Figure A.9	Map showing the overlap of bluefin tuna discards, white marlin discards, and sea turtle interactions for pelagic longline sets from 2001 to 2003. Source: HMS Logbook data 2001-2003			
Figure B.1	Atlantic Albacore Tuna: Spawning, Eggs, and Larvae			
Figure B.2	Atlantic Albacore Tuna: Juvenile B-73			
Figure B.3	Atlantic Albacore Tuna: Adult			
Figure B.4	Atlantic Bigeye Tuna: Spawning, Eggs, and Larvae B-75			
Figure B.5	Atlantic Bigeye Tuna: Juvenile			
Figure B.6	Atlantic Bigeye Tuna: Adult			
Figure B.7	Atlantic Bluefin Tuna: Spawning, Eggs, and Larvae B-78			
Figure B.8	Atlantic Bluefin Tuna: Juveniles B-79			
Figure B.9	Atlantic Bluefin Tuna: Adults			

Figure B.10	Atlantic Skipjack Tuna: Spawning, Eggs, and Larvae	B-81
Figure B.11	Atlantic Skipjack Tuna: Juvenile.	B-82
Figure B.12	Atlantic Skipjack Tuna: Adult	B-83
Figure B.13	Atlantic Yellowfin Tuna: Spawning, Eggs, and Larvae	B-84
Figure B.14	Atlantic Yellowfin Tuna: Juvenile	B-85
Figure B.15	Atlantic Yellowfin Tuna: Adult	B-86
Figure B.16	Atlantic Swordfish: Spawning, Eggs, and Larvae.	B-87
Figure B.17	Atlantic Swordfish: Juvenile	B-88
Figure B.18	Atlantic Swordfish: Adult	B-89
Figure B.19	Blue Marlin: Spawning, Eggs, and Larvae.	B-90
Figure B.20	Blue Marlin: Juvenile	
Figure B.21	Blue Marlin: Adult	B-92
Figure B.22	White Marlin: Spawning, Eggs, and larvae.	B-93
Figure B.23	White Marlin: Juvenile.	B-94
Figure B.24	White Marlin: Adult	B-95
Figure B.25	Sailfish: Spawning, Eggs, and Larvae.	B-96
Figure B.26	Sailfish: Juvenile.	B-97
Figure B.27	Sailfish: Adult.	B-98
Figure B.28	Spearfish: Spawning, Eggs, and Larvae.	B-99
Figure B.29	Spearfish: Juvenile.	B-100
Figure B.30	Spearfish: Adult.	B-101
Figure B.31	Basking Shark: Neonate	B-102
Figure B.32	Basking Shark: Juvenile	B-103
Figure B.33	Basking Shark: Adult.	B-104
Figure B.34	Great Hammerhead: Neonate	B-105
Figure B.35	Great Hammerhead: Juvenile	B-106
Figure B.36	Great Hammerhead: Adult	B-107
Figure B.37	Scalloped Hammerhead: Neonate	B-108
Figure B.38	Scalloped Hammerhead: Juvenile	B-109
Figure B.39	Scalloped Hammerhead: Adult.	B-110
Figure B.40	Smooth Hammerhead: Neonate.	B-111
Figure B.41	Smooth Hammerhead: Juvenile.	B-112
Figure B.42	Smooth Hammerhead: Adult.	B-113
Figure B.43	White Shark: Neonate.	B-114
Figure B.44	White Shark: Juvenile.	B-115
Figure B.45	White Shark: Adult.	B-116
Figure B.46	Nurse Shark: Neonate.	B-117
Figure B.47	Nurse Shark: Juvenile.	B-118
Figure B.48	Nurse Shark: Adult.	B-119
Figure B.49	Bignose Shark: Neonate	B-120
Figure B.50	Bignose Shark: Juvenile	B-121

Figure B.51	Bignose Shark: Adult	B-122
Figure B.52	Blacktip Shark: Neonate.	B-123
Figure B.53	Blacktip Shark: Juvenile.	B-124
Figure B.54	Blacktip Shark: Adult.	B-125
Figure B.55	Bull Shark: Neonate.	B-126
Figure B.56	Bull Shark: Juvenile.	B-127
Figure B.57	Bull Shark: Adult	B-128
Figure B.58	Caribbean Reef Shark: Neonate	B-129
Figure B.59	Caribbean Reef Shark: Juvenile	B-130
Figure B.60	Caribbean Reef Shark: Adult	B-131
Figure B.61	Dusky Shark: Neonate.	B-132
Figure B.62	Dusky Shark: Juvenile.	B-133
Figure B.63	Dusky Shark: Adult	B-134
Figure B.64	Lemon Shark: Neonate.	B-135
Figure B.65	Lemon Shark: Juvenile.	B-136
Figure B.66	Lemon Shark: Adult	B-137
Figure B.67	Night Shark: Neonate	B-138
Figure B.68	Night Shark: Juvenile	B-139
Figure B.69	Night Shark: Adult	B-140
Figure B.70	Sandbar Shark: Neonate	B-141
Figure B.71	Sandbar Shark: Juvenile	B-142
Figure B.72	Sandbar Shark: Adult	B-143
Figure B.73	Sandbar Shark Habitat Area of Particular Concern.	B-144
Figure B.74	Silky Shark: Neonate.	B-145
Figure B.75	Silky Shark: Juvenile.	B-146
Figure B.76	Silky Shark: Adult	B-147
Figure B.77	Spinner Shark: Neonate.	B-148
Figure B.78	Spinner Shark: Juvenile.	B-149
Figure B.79	Spinner Shark: Adult	B-150
Figure B.80	Tiger Shark: Neonate.	B-151
Figure B.81	Tiger Shark: Juvenile.	B-152
Figure B.82	Tiger Shark: Adult.	B-153
Figure B.83	Sand Tiger Shark: Neonate.	B-154
Figure B.84	Sand Tiger Shark: Juvenile.	B-155
Figure B.85	Sand Tiger Shark: Adult.	B-156
Figure B.86	Angel Shark: Neonate	B-157
Figure B.87	Angel Shark: Juvenile	B-158
Figure B.88	Angel Shark: Adult	B-159
Figure B.89	Bonnethead Shark : Neonate	B-160
Figure B.90	Bonnethead Shark: Juvenile	B-161
Figure B.91	Bonnethead Shark: Adult	B-162

Figure B.92	Altantic Sharpnose: Neonate	. B-163
Figure B.93	Atlantic Sharpnose: Juvenile	. B-164
Figure B.94	Atlantic Sharpnose Shark: Adult.	. B-165
Figure B.95	Blacknose Shark: Neonoate.	. B-166
Figure B.96	Blacknose Shark: Juvenile.	. B-167
Figure B.97	Blacknose Shark: Adult.	. B-168
Figure B.98	Finetooth Shark: Neonate.	. B-169
Figure B.99	Finetooth Shark: Juvenile.	. B-170
Figure B.100	Finetooth Shark: Adult	. B-171
Figure B.101	Smalltail Shark: All Life Stages.	. B-172
Figure B.102	Bigeye Sixgill Shark: All Life Stages.	. В-173
Figure B.103	Sevengill Shark: All Life Stages	. B-174
Figure B.104	Sixgill Shark: All Life Stages.	. B-175
Figure B.105	Longfin Mako Shark: Neonate	. B-176
Figure B.106	Longfin Mako Shark: Juvenile	. В-177
Figure B.107	Longfin Mako Shark: Adult	. B-178
Figure B.108	Porbeagle Shark: Neonate.	. B-179
Figure B.109	Porbeagle Shark: Juvenile.	. B-180
Figure B.110	Porbeagle Shark: Adult.	. B-181
Figure B.111	Shortfin Mako Shark: Neonate.	. B-182
Figure B.112	Shortfin Mako Shark: Juvenile.	. B-183
Figure B.113	Shortfin Mako Shark: Adult	. B-184
Figure B.114	Blue Shark: Neonate.	. B-185
Figure B.115	Blue Shark: Juvenile.	. B-186
Figure B.116	Blue Shark: Adult.	. B-187
Figure B.117	Oceanic Whitetip Shark: Neonate	. B-188
Figure B.118	Oceanic Whitetip Shark: Juvenile	. B-189
Figure B.119	Oceanic Whitetip Shark: Adult	. B-190
Figure B.120	Bigeye Thresher Shark: Juvenile.	. B-191
Figure B.121	Bigeye Thresher Shark: Adult.	. B-192
Figure B.122	Thresher Shark: Neonate	. B-193
Figure B.123	Thresher Shark: Juvenile	. B-194
Figure B.124	Thresher Shark: Adult	. B-195

LIST OF COMMONLY USED ABBREVIATIONS AND ACRONYMS

AA	Assistant Administrator for Fisheries
ACCSP	Atlantic Coastal Cooperative Statistics Program
ACS	Angler consumer surplus
ALRS	Automated Landings Reporting System
ALWTRP	Atlantic Large Whale Take Reduction Plan
ALWTRT	Atlantic Large Whale Take Reduction Team
ANPR	Advanced Notice of Proposed Rulemaking
AOCTRP	Atlantic Offshore Cetacean Take Reduction Plan
AOCTRT	Atlantic Offshore Cetacean Take Reduction Team
AP	Advisory Panel
APA	Administrative Procedure Act
ASMFC	Atlantic States Marine Fisheries Commission
ATCA	Atlantic Tunas Convention Act
В	Biomass
BAYS	Bigeye, albacore, yellowfin, skipjack tunas
BET	Bigeye tuna
ВЕТҮР	Bigeye Tuna Year Program
BFT	Bluefin tuna
BiOp	Biological Opinion
BLL	Bottom Longline
B_{MSY}	Biomass expected to yield maximum sustainable yield
B_{OY}	Biomass expected to yield optimum yield
BSD	Bluefin Tuna Statistical Document
BTF	By the fish
BUM	Blue marlin
CAR	Caribbean Statistical Area
СВР	Customs and Border Protection
CFDBS	Commercial Fisheries Database System
CFMC	Caribbean Fishery Management Council
CFL	Curved fork length
CFR	Code of Federal Regulations
СНВ	Charter/Headboat
CIAT	Spanish for IATTC
CIE	Center for Independent Experts
CITES	Convention on International Trade in Endangered Species of Wild Fauna and Flora
COE	Certificate of Eligibility
COFI	Committee on Fisheries
СРІ	Consumer Price Index

CPUE	Catch per unit effort
CSFOP	Commercial Shark Fishery Observer Program (run by University of Florida)
CSR	Center for Shark Research
CSTP	Cooperative Shark Tagging Program
CZMA	Coastal Zone Management Act
DEIS	Draft Environmental Impact Statement
DPS	Distinct population segment
DRG	Dredge
DSGFOP	
	Directed Shark Gillnet Fishery Observer Program Dressed weight
dw EA	Environmental Assessment
EEZ	Exclusive Economic Zone
EFC	East Florida Coast closed area
EFH	Essential fish habitat
EFP	Exempted fishing permit
EIS	Environmental Impact Statement
ЕО	Executive Order
ESA	Endangered Species Act
F	Instantaneous fishing mortality
FAD	Fish aggregating device
FAO	Food and Agriculture Organization
FAS	Free Alongside Ship
FEC	Florida East Coast Statistical Area
FEIS	Final Environmental Impact Statement
FL	Fork length
FMP	Fishery management plan
F_{MSY}	Instantaneous fishing mortality rate expected to yield maximum sustainable yield
FMU	Fishery management unit
F_{OY}	Fishing mortality rate expected to yield optimum yield
FR	Federal Register
FRFA	Final regulatory flexibility analysis
GDP	Gross Domestic Product
GIS	Geographic Information System
GOM	Gulf of Mexico
GSAFDF	Gulf and South Atlantic Fishery Development Foundation
GMFMC	Gulf of Mexico Fishery Management Council
GSMFC	Gulf States Marine Fisheries Commission
НАССР	Hazard Analysis Critical Control Point
HAPC	Habitat area of particular concern
I	1

HBS	Headboat Survey
HMS	Highly migratory species: Atlantic sharks, tunas, swordfish, and billfish
HTS	Harmonized Tariff Schedule
IATTC	Inter-American Tropical Tuna Commission
ICCAT	International Commission for the Conservation of Atlantic Tunas
ILAP	Initial limited access permit
IMARPE	Instituo del Mar del Peru
INP	Instituto Nacional de Pesca
IPOA	International Plan of Action
IRFA	Initial regulatory flexibility analysis
ITP	International Trade Permit
ITQ	Individual transferable quota
ITS	Incidental take statement
IUU	Illegal, Unregulated, and Unreported
LAP	Limited access permit
LCS	Large coastal sharks
LJFL	Lower jaw fork length
LOA	Letter of Acknowledgment
LOF	List of Fisheries
LPS	Large Pelagic Survey
LWTRP	Large Whale Take Reduction Plan
LWTRT	Large Whale Take Reduction Team
M	Mortality
MAB	Mid-Atlantic Bight Statistical Area
MAFMC	Mid-Atlantic Fishery Management Council
Magnuson-Stevens Act	Magnuson-Stevens Fishery Conservation and Management Act
MFMT	Maximum fishing mortality threshold
MMPA	Marine Mammal Protection Act
MMS	Minerals Management Service
MPA	Marine protected area
MRFSS	Marine Recreational Fishing Statistics Survey
MSST	Minimum stock size threshold
MSY	Maximum sustainable yield
mt	Metric tons
NCA	North Central Atlantic
NEC	Northeast Coastal Statistical Area
NED	Northeast Distant Statistical Area
NEFMC	New England Fishery Management Council
NEFSC	Northeast Fisheries Science Center, NMFS

NEPA	National Environmental Policy Act
NERO	Northeast Regional Office, NMFS
NFRDI	National Fisheries Research and Development Institute
NGO	Non-governmental organization
NMFS	National Marine Fisheries Service
nmi	Nautical mile
NOA	Notice of Availability
NOAA	National Oceanographic and Atmospheric Administration
NOI	Notice of Intent
NPOA	National Plan of Action
NRC	Natural Resources Consultants, Inc.
NS	National Standards
NYB	New York Bight
OSF	Office of Sustainable Fisheries
OY	Optimum yield
PAT	Pop-up archival tag
PFD	Personal flotation device
PIFSC	Pacific Islands Fisheries Science Center
PLL	Pelagic longline
PLTRP	Pelagic Longline Take Reduction Plan
PLTRT	Pelagic Longline Take Reduction Team
POP	Pelagic observer program
PPI	Producer price index
OPR	Office of Protected Resources
PRA	Paperwork Reduction Act
PRM	Post-release mortality
PSAT	Pop-up satellite archival tag
RBS	Recreational Billfish Survey
Reg Flex Act	Regulatory Flexibility Act
RIR	Regulatory Impact Review
RFMC	Regional Fishery Management Council
RPAs	Reasonable and Prudent Alternatives
RPMs	Reasonable and Prudent Measures
RUM	Random utility model
SAFE Report	Stock Assessment and Fishery Evaluation Report
SAFMC	South Atlantic Fishery Management Council
SAB	South Atlantic Bight
SAI	Sailfish
SAR	Sargasso Sea

SBR	Spawning Stock Biomass Ratio
SCRS	Standing Committee for Research and Statistics
SCS	Small coastal sharks
SCUBA	Self contained underwater breathing apparatus
SD	Statistical document
Secretary	Secretary of Commerce
SEDAR	Southeast Data, Assessment, and Review
SEFSC	Southeast Fisheries Science Center, NMFS
SEIS	Supplemental Environmental Impact Statement
SEN	Seines
SERO	Southeast Regional Office, NMFS
SEW	Stock evaluation workshop
SFA	Sustainable Fisheries Act
SFL	Straight fork length
SK Program	Saltonstall-Kennedy Program
SRP	Scientific research permit
SSB	Spawning stock biomass
SWFSC	Southwest Fisheries Science Center
TAC	Total allowable catch
TAG	Tag-A-Giant
TAL	Total allowable landings
TCs	Terms and Conditions
TL	Total length
TRP	Traps and pots
TUNS	Tuna North and Tuna South
TWL	Trawls
TXPWD	Texas Parks and Wildlife Department
UNK	Unknown
USFWS	United States Fish and Wildlife Service
VIMS	Virginia Institute of Marine Science
VMS	Vessel monitoring system
WHM	White marlin
WPFMC	Western Pacific Fishery Management Council
WTP	Willingness to pay
ww	Whole weight
WWF	World Wildlife Fund
YFT	Yellowfin tuna
YOY	Young of the year